

# MATH 595 (Group Cohomology) Notes

Jiantong Liu

November 13, 2023

## 1 AUG 21, 2023: INTRODUCTION

Group cohomology works over different settings of groups, like finite groups, profinite groups, and topological groups. The course will develop towards

- duality in  $H^*(G, -)$ , and
- focus on computations, e.g., spectral sequences.

We first establish some notations.

- Let  $G$  be a group. If  $G$  has a topology, that would also be part of the information of  $G$ .
- A (left)  $G$ -module is an abelian group  $M$  with an action map

$$\begin{aligned} G \times M &\rightarrow M \\ (g, m) &\mapsto g \cdot m = gm \end{aligned}$$

satisfying

- $1 \cdot m = m$ ,
- $(gh) \cdot m = g \cdot (hm)$ ,
- $g(m + m') = gm + gm'$ .

**Remark 1.1.** If  $G$  is a finite group, then the associated (non-commutative) group ring  $\mathbb{Z}[G] := \bigoplus_{g \in G} \mathbb{Z}e_g$ , where the multiplication is determined by  $e_g e_h = e_{gh}$ . Therefore, a  $G$ -module is just a  $\mathbb{Z}[G]$ -module.

**Example 1.2.** • Trivial module  $\mathbb{Z}$ , or any abelian group with the trivial action  $g \cdot a = a$ .

- $C_2$ , or any group with  $f : G \twoheadrightarrow C_2$ , then  $G$  with  $C_2$  as a quotient gives the sign representation  $\mathbb{Z}_{\text{sgn}}$ , with  $g \cdot (a) = (-1)^{\rho(g)} a$ .
- $\mathbb{Z}[G]$  is a  $G$ -module via the left multiplication action, and/or the conjugation action.

**Definition 1.3** (Fixed points/Invariants). The set of fixed points of  $M$  over  $G$  is  $M^G = \{m \in M \mid gm = m \ \forall g \in G\}$ .

**Definition 1.4** (Orbits/Coinvariants). The set of orbits of  $M$  over  $G$  is  $M_G = M/(gm - m)$ .

**Example 1.5.** If  $M = \mathbb{Z}_{\text{sgn}}$ , then everything gets multiplied by  $-1$ , so there are no fixed points. The orbits of  $M$  over  $G$  would be  $\mathbb{Z}_{\text{sgn}}/(-2) \cong \mathbb{Z}/2\mathbb{Z}$ .

**Example 1.6.** If  $M = \mathbb{Z}[G]$ , then the fixed points are  $\mathbb{Z} \left\{ \sum_{g \in G} e_g \right\}$ .

Thinking in a categorical setting, we have a trivial action function  $\mathbb{Z}\text{-Mod} \rightarrow G\text{-Mod}$ , sending  $ga \mapsto a$  for all  $g \in G$  and  $a \in A$ . This gives an exact functor from  $\mathbf{Ab}$  to  $G\text{-Mod}$ . Then this functor has a right adjoint  $( )^G : G\text{-Mod} \rightarrow \mathbf{Ab}$ , and a left adjoint  $( )_G : \mathbf{Ab} \rightarrow G\text{-Mod}$ . More specifically,  $M^G$  becomes the maximal trivial action submodule of  $M$ , namely  $\text{Hom}_G(\mathbb{Z}, M)$ ;  $M_G$  becomes the largest quotient of  $M$  with trivial action, namely  $\mathbb{Z} \otimes_{\mathbb{Z}[G]} M$ . This simplifies to the tensor-hom adjunction in some sense. For a more detailed derivation of this, see Chapter 6.1 of Weibel.

**Remark 1.7.** In general, as in the category of  $G$ -sets, we have the orbit functor  $X \mapsto X/G$  and the fixed point functor  $X \mapsto X^G$ . The orbit functor is left adjoint to the free  $G$ -set functor, and the fixed point functor is the right adjoint of the trivial  $G$ -set functor.

**Remark 1.8.** Read more about the setting in profinite groups with their topologies in Neukirch-Schmidt-Wingberg.

**Definition 1.9** (Profinite Group). A profinite group of a collection of groups is  $G = \varprojlim_i G_i$  as an inverse limit, where each  $G_i$  is a finite group of the form  $G/U_i$  for some open  $U_i$ . This gives a topology to the profinite group.

**Remark 1.10.** The groups rings  $\mathbb{Z}[[G]] = \varprojlim_i \mathbb{Z}[G_i]$ . For instance, let  $G = \hat{\mathbb{Z}}_p = \varprojlim_n \mathbb{Z}/p^n\mathbb{Z}$ , then  $\mathbb{Z}_p[[G]] = \varprojlim_n \mathbb{Z}_p[\mathbb{Z}/p^n\mathbb{Z}]$ , where each  $\mathbb{Z}[\mathbb{Z}/p^n\mathbb{Z}] \cong \bigoplus_{i=0}^{n-1} \mathbb{Z}\{e_i\}$  where  $e_i \cdot e_j = e_{ij}$ . Therefore,  $\mathbb{Z}_p[[G]]$  is now equivalent to  $\varprojlim_n \mathbb{Z}_p[t]/(t^{p^n} - 1_e)$ , and hence becomes a power series.

**Remark 1.11.** By a change of variables, this becomes  $\varprojlim_n \mathbb{Z}_p[x]/(x^{p^n})$ , but this only works in the finite group  $\mathbb{Z}_p$  case, and not in general for  $\mathbb{Z}$ .

**Example 1.12.**  $\mathbb{Z}[C_n] \cong \mathbb{Z}\{e\} \oplus \mathbb{Z}\{g\} \oplus \mathbb{Z}\{g^2\} \oplus \cdots \oplus \mathbb{Z}\{g^{n-1}\} \cong \mathbb{Z}[g]/(g^n - 1_e)$ .

## 2 AUG 23, 2023: COHOMOLOGY OF GROUPS

**Definition 2.1.** Let  $G$  be a group, then we have a diagram

$$EG : \cdots \rightrightarrows G \times G \rightrightarrows G$$

where the arrows are given by

$$EG^n = G^{n+1} \xrightarrow{d_i} G^n$$

for all  $0 \leq i \leq n$ . In the sense of simplicial sets, we have  $d_i(g_0, \dots, g_n) = (g_0, \dots, \hat{g}_i, \dots, g_n)$ .

Now let  $M$  be a  $G$ -module, then we define  $X^n = X^n(G, M) = \text{Map}_{\text{Set}}(G^{n+1}, M)$ .  $G$  now has an action on this set, given by

$$(g \circ f)(g_0, \dots, g_n) = gf(g^{-1}g_0, \dots, g^{-1}g_n).$$

The action on  $d^i$ 's are contravariant, namely we obtain  $d_i^* : X_n \rightarrow X^{n+1}$  with an inherited structure. Note that  $M$  sits inside  $X^0$ , therefore we have a complex  $(*)$ :

$$0 \longrightarrow M \xleftarrow{\partial_0} X^0 \xrightarrow{\partial_1} X^1 \xrightarrow{\partial_2} X^2 \xrightarrow{\partial_3} \cdots$$

Here  $\partial_0$  includes  $M$  as the constant functions into  $X$ , namely  $\partial_0(m) = f$  for  $f(g) = m$ , and so on. In general, for  $n > 0$ , we have

$$\partial_n = \sum_{i=0}^n (-1)^i d_i^*.$$

**Lemma 2.2.** The complex  $(*) : M \rightarrow X^\cdot$  is an exact complex of  $G$ -modules, i.e.,  $\partial^2 = 0$  and  $\ker(\partial_{n+1}) = \text{im}(\partial_n)$ , and the  $\partial_i$ 's preserves the  $G$ -action. This is called the standard resolution of  $M$  as a  $G$ -module.

*Proof.* Exercise. □

**Definition 2.3.** The  $G$ -fixed points of the  $X^n$ 's are defined by  $C^n(G, M) = (X^n(G, M))^G$ , called the homogeneous  $n$ -cochains of  $G$  with coefficients in  $M$ . Because the complex preserves  $G$ -actions, then we obtain a complex of  $C^n(G, M)$ 's, given by

$$0 \longrightarrow C^0(G, M) \xrightarrow{\partial_0} C^1(G, M) \xrightarrow{\partial_1} \dots$$

**Remark 2.4.** To see what the induced mapping is, suppose  $A \rightarrow B$  is a  $G$ -module map, then there is an induced map of fixed points  $A^G \rightarrow B^G$  by the restriction. In particular, let  $a \in A$  be fixed with  $ga = a$  for all  $g \in G$ , then  $f(a) = f(ga) = gf(a)$ .

**Remark 2.5.** In the complex of Definition 2.3,  $\partial^2 = 0$  as well, but in general this is not an exact sequence.

**Definition 2.6** (Group Cohomology). The group cohomology of  $G$  with coefficients in  $M$  is the collection

$$\{H^n(G, M)\}_{n \geq 0},$$

where  $H^n(G, M) := H^n(C^\cdot(G, M)) = \ker(\partial : C^n \rightarrow C^{n+1}) / \text{im}(\partial : C^{n-1} \rightarrow C^n)$ . We usually use the notion of cocycles  $Z^n(G, M) = \ker(\partial : C^n \rightarrow C^{n+1})$  and coboundaries  $B^n(G, M) = \text{im}(\partial : C^{n-1} \rightarrow C^n)$ .

**Exercise 2.7.** Show that  $H^0(G, M)$  is isomorphic to  $M^G$ .

**Definition 2.8.** The inhomogeneous cochains  $C_i^\cdot(G, M)$  are given by

- $C_i^0 = M$ , and
- for  $n > 0$ ,  $C_i^n = \text{Map}(G^n, M)$ ,

with coboundary maps  $\partial^{n+1} : C_i^n \rightarrow C_i^{n+1}$ , given by

- $\partial^1(m)(g) = gm - m$ ,
- $\partial^2(f)(g_1, g_2) = g_1 f(g_2) - f(g_1 g_2) + f(g_1)$ , and so on, with
- $\partial^{n+1}(f)(g_1, \dots, g_{n+1}) = g_1 f(g_2, \dots, g_{n+1}) + \sum_{i=1}^n (-1)^i f(g_1, \dots, g_i g_{i+1}, \dots, g_{n+1}) + (-1)^{n+1} f(g_1, \dots, g_n)$ .

This gives the inhomogeneous setting of this cochain.

**Lemma 2.9.** The maps

$$\begin{aligned} C^n(G, M) &\rightarrow C_i^n(G, M) \\ (\varphi : G^{n+1} \rightarrow M) &\mapsto (f : G^n \rightarrow M) \\ f(g_1, \dots, g_n) &:= \varphi(1, g_1, g_1 g_2, \dots, g_1 g_2 \cdots g_n) \end{aligned}$$

give a cochain homotopy equivalence  $C^\cdot(G, M) \xrightarrow{\sim} C_i^\cdot(G, M)$ , and hence this is a quasi-isomorphism.

**Corollary 2.10.** The cohomology  $H^*(C_i^\cdot(G, M)) \cong H^*(G, M)$ .

**Remark 2.11.** Any cohomology class can be represented by a normalized inhomogeneous cocycle  $f : G^n \rightarrow M$ , i.e.,  $f(g_1, \dots, g_n) = 0$  where  $g_i = 1$  for some  $i$ .

**Remark 2.12.** Even for  $G = C_2$ ,  $C_i^n$  or  $C^n$  get large as  $n$  grows.

**Remark 2.13.** • Using homological algebra, we can find other cochain complexes which computes group cohomology  $H^*(G, M)$ .

- We would also understand  $H^*(G, M)$  as the failure of exactness of  $(\ )^G : G\text{-Mod} \rightarrow \mathbf{Ab}$ . Therefore, when taking the fixed points, the exact sequence may not be mapped to another exact sequence. In particular, if we take an exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

of  $G$ -modules, the induced sequence

$$0 \longrightarrow A^G \longrightarrow B^G \longrightarrow C^G$$

do not give a surjection at  $B^G \rightarrow C^G$ . One needs to take higher cohomology to obtain a long exact sequence. Hence,  $(\ )^G : G\text{-Mod} \rightarrow \mathbf{Ab}$  is a left exact functor, but not necessarily right exact.

## 3 AUG 25, 2023: COHOMOLOGY OF GROUPS, CONTINUED

**Example 3.1.** Let  $G$  be  $C_2$ , or any group with a surjection  $p$  onto  $C_2$ , then it has an action on  $\mathbb{Z}_{\text{sgn}}$  given by  $g \cdot a = (-1)^{p(g)}a$ , therefore we have a short exact sequence

$$0 \longrightarrow \mathbb{Z}_{\text{sgn}} \xrightarrow{\times 2} \mathbb{Z}_{\text{sgn}} \longrightarrow \mathbb{Z}/2\mathbb{Z} \longrightarrow 0$$

and taking the fixed point functor we have

$$0 \longrightarrow 0 \longrightarrow 0 \longrightarrow \mathbb{Z}/2\mathbb{Z}.$$

**Remark 3.2.** Higher homologies measure the failure of exactness.

**Remark 3.3.** The collection  $\{H^n(G, -)\}_{n \in \mathbb{Z}}$  satisfies

- $H^n(G, -) = 0$  for  $n < 0$ ;
- for short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  in  $G\text{-Mod}$ , we have a long exact sequence

$$0 \longrightarrow H^0(G, A) \longrightarrow H^1(G, B) \longrightarrow H^1(G, C) \xrightarrow{\delta} H^1(G, A) \longrightarrow \cdots$$

where  $\delta$  is the connecting homomorphism.

- the connecting homomorphisms  $\delta$  are natural, i.e., given a commuting diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & B & \longrightarrow & C \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & A' & \longrightarrow & B' & \longrightarrow & C' \longrightarrow 0 \end{array}$$

the induced diagram

$$\begin{array}{ccc} H^n(G, C) & \xrightarrow{\delta} & H^{n+1}(G, A) \\ \downarrow & & \downarrow \\ H^n(G, C') & \xrightarrow{\delta} & H^{n+1}(G, A') \end{array}$$

also commutes, and  $\{H^n(G, -)\}_{n \in \mathbb{Z}}$  is a cohomological  $\delta$ -functor. Note that a  $\delta$ -functor is additive, and usually occurs in abelian categories.

**Definition 3.4** ( $\delta$ -functor). A map of  $\delta$ -functors  $T^* \rightarrow F^*$  is a collection of natural transformations  $T^n \rightarrow F^n$ , commuting with the  $\delta$ 's, i.e.,

$$\begin{array}{ccc} T^n & \longrightarrow & F^n \\ \delta_T \downarrow & & \downarrow \delta_F \\ T^{n+1} & \longrightarrow & F^{n+1} \end{array}$$

A  $\delta$ -functor  $T^*$  is universal if, given any other  $\delta$ -functor  $F^*$ , a map  $T^* \rightarrow F^*$  is uniquely determined by  $T^0 \rightarrow F^0$ .

**Proposition 3.5.**  $H^*(G, -) : G\text{-Mod} \rightarrow \mathbf{Ab}$  is a  $\delta$ -functor.

*Proof.* We need to show:

- each  $H^n(G, -)$  is a well-defined functor,
- the connecting homomorphisms  $\delta$ 's gives a long exact sequence,
- the naturality of  $\delta$ .

First, let  $f : A \rightarrow B$  be in  $G\text{-Mod}$ , then  $C^*(G, A) \rightarrow C^*(G, B)$  is equivalent to  $\text{Map}(G^{*+1}, A)^G \rightarrow \text{Map}(G^{*+1}, B)^G$  by composition with  $f$ . One can show that this is equivariant, i.e., respects the  $G$ -action, so it is well-defined to take the fixed points, and thus commutes with  $\partial$ 's.

Second, we need to apply the snake lemma. Given a short exact sequence  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ , we claim:

**Claim 3.6.**  $0 \longrightarrow C^*(G, A) \longrightarrow C^*(G, B) \longrightarrow C^*(G, C) \longrightarrow 0$  is a short exact sequence of cochain complexes, i.e.,  $C^*(G, -) : G\text{-Mod} \rightarrow \mathbf{coCh}$  is an exact functor.

*Subproof.* Exercise. ■

Now take the complex

$$\begin{array}{ccccccc} 0 & \longrightarrow & C^n(G, A) & \longrightarrow & C^n(G, B) & \longrightarrow & C^n(G, C) \longrightarrow 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & C^{n+1}(G, A) & \longrightarrow & C^{n+1}(G, B) & \longrightarrow & C^{n+1}(G, C) \longrightarrow 0 \end{array}$$

and quotient the boundaries everywhere (and thus lose the injectivity/surjectivity when applicable)

$$\begin{array}{ccccccc} C^n(G, A)/B^n(G, A) & \longrightarrow & C^n(G, B)/B^n(G, B) & \longrightarrow & C^n(G, C)/B^n(G, C) & \longrightarrow & 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & Z^{n+1}(G, A) & \longrightarrow & Z^{n+1}(G, B) & \longrightarrow & Z^{n+1}(G, C) \end{array}$$

Taking the kernels and cokernels on  $\partial$ 's, we obtain a complex

$$\begin{array}{ccccccc} & & 0 & & 0 & & 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ & & H^n(G, A) & \longrightarrow & H^n(G, B) & \longrightarrow & H^n(G, C) \\ & & \downarrow & & \downarrow & & \downarrow \\ C^n(G, A)/B^n(G, A) & \longrightarrow & C^n(G, B)/B^n(G, B) & \longrightarrow & C^n(G, C)/B^n(G, C) & \longrightarrow & 0 \\ & & \downarrow \partial & & \downarrow \partial & & \downarrow \partial \\ 0 & \longrightarrow & Z^{n+1}(G, A) & \longrightarrow & Z^{n+1}(G, B) & \longrightarrow & Z^{n+1}(G, C) \\ & & \downarrow & & \downarrow & & \downarrow \\ & & H^{n+1}(G, A) & \longrightarrow & H^{n+1}(G, B) & \longrightarrow & H^{n+1}(G, C) \end{array}$$

By the snake lemma, we obtain the long exact sequence. □

**Proposition 3.7.** If  $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$  is a short exact sequence such that  $H^*(G, B) = 0$  for  $* > 0$  (or at least  $H^n(G, B) = 0 = H^{n+1}(G, B)$ ), then  $\delta : H^n(G, C) \rightarrow H^{n+1}(G, A)$  is an isomorphism.

**Definition 3.8** (Acyclic, Cohomologically Trivial). A  $G$ -module  $M$  is

- acyclic if  $H^*(G, M) = 0$  for  $* > 0$ ,
- cohomologically trivial if  $H^*(H, M) = 0$  for  $* > 0$  and any (closed) subgroup  $H \subseteq G$ .

**Definition 3.9** (Induced Module). Given any  $G$ -module  $M$ , the induced module  $\text{ind}_G(M) = \text{Map}(G, M) = X^0(G, M)$ .

**Example 3.10.**  $M$  could have the trivial action.

**Exercise 3.11.** For any  $M$ , the induced module of  $M$  over  $G$  is isomorphic (under the  $G$ -action) to the induced module of module given by forgetful action over  $G$ .

**Remark 3.12.** •  $\text{Ind}_G(-) : G\text{-Mod} \rightarrow G\text{-Mod}$  is exact.

- We say  $A$  is an induced module if  $A \cong \text{Ind}_G(M)$  for some module  $M$ . If  $A$  is an induced  $G$ -module, then  $A$  is induced as an  $H$ -module for any subgroup  $H \subseteq G$ .

**Lemma 3.13.** Induced modules are cohomologically trivial.

*Proof.* There is an isomorphism

$$C^*(G, \text{Ind}_G(M)) \cong X^*(G, M).$$

□

**Remark 3.14.** We have an equivariant inclusion of fixed points

$$M \hookrightarrow \text{Ind}_G(M)$$

which is an embedding, and we take  $Q \cong \text{Ind}_G(M)/M$ , then this extends to a short exact sequence

$$0 \longrightarrow M \hookrightarrow \text{Ind}_G(M) \longrightarrow Q \longrightarrow 0$$

then  $H^{n+1}(G, M) \cong H^n(G, Q)$ . One say that  $H^*(G, -)$  is effaceable. By Tohoku, an effaceable is universal.

#### 4 AUG 28, 2023: FIRST COHOMOLOGY OF GROUPS

There are three ways to think about  $H^1(G, M)$ .

##### 4.1 CROSSED HOMOMORPHISMS

Recall that  $H^1(G, M) = Z_i^1(G, M)/B_i^1(G, M)$  as inhomogeneous cochains, where

- $Z_i^1(G, M) = \ker(\text{Map}(G, M) \rightarrow \text{Map}(G \times G, M))$  where the map sends  $f \mapsto (g, h) \mapsto gf(h) - f(gh) + f(g)$ . The kernel of this is exactly the maps  $f$  such that  $f(gh) = gf(h) + f(g)$ , and note that this is not a group homomorphism.
- $B_i^1(G, M) = \text{im}(M \rightarrow \text{Map}(G, M))$  given by  $m \mapsto (g \mapsto gm - m)$ , where the image is called a principal crossed homomorphism.

**Exercise 4.1.**  $B_i^1(G, M) \cong M/M^G$  as an isomorphism of  $\mathbb{Z}[G]$ -modules.

**Remark 4.2.** If the  $G$ -action is trivial, then  $H^1(G, M) = \text{Hom}_{\text{Grp}}(G, M)$ .

**Corollary 4.3.** If  $G$  is a finite group with trivial action, then  $H^1(G, \mathbb{Z}) = 0$ .

**Theorem 4.4** (Hilbert's Theorem 90). Let  $L/K$  be a Galois extension with (finite or profinite) Galois group  $G$ , then  $H^1(G, L^\times) = 0$ .

*Proof.* Let  $f : G \rightarrow L^\times$  be a crossed homomorphism. We know the addition is given by  $f(gh) = gf(h) + f(g)$ , and the multiplication is given by  $f(gh) = (g \cdot f(h))f(g)$ , where  $\cdot$  represents the group action. Now for any  $l \in L^\times$ , the multiplication with respect to  $l$  is given by  $m_l = \sum_{h \in G} f(h)(h \cdot l)$ . We can first choose  $l$  so that  $m_l \neq 0$ , since the Galois conjugates  $h \cdot l$  over  $l \in L$  are linearly independent. For  $g \in G$ , we have

$$\begin{aligned} g \cdot m_l &= \sum_{h \in G} (g \cdot f(h))(gh \cdot l) \\ &= \sum_{h \in G} \frac{f(gh)}{f(g)} (gh \cdot l) \\ &= \frac{1}{f(g)} \sum_{h \in G} f(gh)(gh \cdot l) \\ &= \frac{1}{f(g)} m_l. \end{aligned}$$

Therefore,  $f(g) = \frac{m_l}{g \cdot m_l}$ . For any crossed homomorphism, there exists  $m \in L^\times$  such that  $f(g) = \frac{gm}{m}$ , so every crossed homomorphism is principal. □

**Exercise 4.5.** Let  $G$  acts over a commutative ring  $R$ , then  $H^1(G, R^\times)$  classifies invariant  $R$ -modules with a compatible  $G$ -action.

4.2 NON-ABELIAN  $H^1$  AND TORSORS

Let  $A$  be a group with  $G$ -action, so let the action  $g \cdot a = {}^g a$ . Hence,  $g \cdot (ab) = {}^g a {}^g b$ . Define the  $G$ -cocycles to be  $f : G \rightarrow A$  such that  $f(gh) = f(g) {}^g f(h)$ . Two cocycles  $f$  and  $f'$  are said to be cohomologous as  $f \sim f'$  if there exists  $a \in A$  such that for all  $g \in G$ ,  $f'(g) = a^{-1} f(g) {}^g a$ . This becomes an equivalence relation on the set of  $G$ -cocycles with coefficients in  $A$ , then  $H^1(G, A)$  is the set of equivalence classes of  $G$ -cocycles. Now the first cohomology  $H^1(G, A)$  has only a pointed set structure with distinguished point  $f \equiv 1$ , the constant function at 1.

**Exercise 4.6.** This definition is equivalent to the inhomogeneous cochain definition in the abelian case.

**Definition 4.7.** An  $A$ -torsor is a  $G$ -set  $X$  with action

$$\begin{aligned} X \times A &\rightarrow A \\ (x, a) &\mapsto xa \end{aligned}$$

that is free and transitive, i.e., for any  $x, y \in X$ , there exists a unique  $a \in A$  such that  $y = xa$ . Moreover, the action  $X \times A \rightarrow X$  respects the  $G$ -action, i.e.,  ${}^g(xa) = {}^g x {}^g a$ .

**Remark 4.8.** •  $A$  is an  $A$ -torsor.

- An isomorphism of  $A$ -torsors is a bijection that respects the  $G$ - and  $A$ - action.
- If  $A \subseteq B$  is a sub- $G$ -group, then  $bA$  is an  $A$ -torsor.
- An  $A$ -torsor is a principal  $A$ -bundle on the classifying space  $BG$ .

**Theorem 4.9.** There is a canonical bijection of pointed sets

$$H^1(G, A) \cong \text{Torsor}(G, A)$$

*Proof.* • The backwards map  $\lambda : \text{Torsor}(G, A) \rightarrow H^1(G, A)$  is defined as follows: for  $x \in \text{Torsor}(G, A)$ , we want to define a cocycle  $f(X) : G \rightarrow A$ . For arbitrary  $x \in X$ , note that for any  $g \in G$ , there exists a unique  $f_x(g) \in A$  such that  ${}^g x = x f_x(g)$  by the simple transitivity of the  $A$ -action on  $X$ . To see this is well-defined, if we have another  $y \in X$ , then  $y = xb$  for some  $b \in A$ , then  $f_y(g) = b^{-1} f_x(g) {}^g b$ , so  $f_x$  and  $f_y$  are cohomologous and define the same class in  $H^1(G, A)$ , which is defined to be the image  $\lambda(X)$ .

- To define  $\mu : H^1(G, A) \rightarrow \text{Torsor}(G, A)$ , given a cocycle  $f : G \rightarrow A$ , let  $X_f$  be the group  $A$ , then the action of  $A$  on  $X_f$  is by multiplication on the right, and one can twist the  $G$ -action on it using cocycle  $f : G \rightarrow A$  with  ${}^g x = f(g)gx$ , which defines an  $A$ -torsor. This is well-defined.

□

**Remark 4.10.** Suppose

$$1 \longrightarrow A \longrightarrow B \xrightarrow{p} C \longrightarrow 1$$

is a short exact sequence of  $G$ -groups, i.e.,  $A$  is a sub- $G$ -group and  $C \cong B/A$ , then there is a long exact sequence

$$1 \longrightarrow A^G \longrightarrow B^G \longrightarrow C^G \xrightarrow{\delta} H^1(G, A) \longrightarrow H^1(G, B) \longrightarrow H^1(G, C)$$

where  $\delta$  is given by  $\delta(c) = p^{-1}(c)$ . For the exactness in the sense of pointed sets to work, the kernel is the subset mapping to the distinguished element.

## 4.3 EXTENSION SPLITTING

Consider the a split extension

$$1 \longrightarrow A \longrightarrow E \xrightarrow{p} G \longrightarrow 1$$

That is,  $E$  is the direct product  $A \times G$  with group action  $(a, g)(a', g') = (a {}^g a', gg')$ , and by definition  $E$  is the semidirect product  $A \rtimes G$ . Equivalently, there exists a section (as group homomorphism)  $s : G \rightarrow E$ .

There is an equivalence relation on the set of sections to the projection  $p : E \rightarrow G$ , where the sections  $s, s' : G \rightarrow E$  are conjugates if there exists  $a \in A$  such that  $s'(g) = a^{-1} s(g) a$ . We denote  $\text{sec}(E \rightarrow G)$  to be the conjugacy class of sections of  $p$ . Note that the class of trivial section  $s : g \mapsto (1, g) \in E$  is the distinguished element.

**Proposition 4.11.** The pointed set  $H^1(G, A)$  is isomorphic to  $\text{sec}(E \rightarrow G)$ .

*Proof.* Take  $\varphi \in \text{sec}(E \rightarrow G)$ , then the composition  $G \xrightarrow{\varphi} E \xrightarrow{\pi_1} A$ , where  $\pi_1$  is the set-theoretic projection to the first component, defines a cocycle  $G \rightarrow A$ . Conversely, given a cocycle  $f : G \rightarrow A$ , the section is given by  $g \mapsto (f(g), g)$ .  $\square$

**Exercise 4.12.** Expand the proof above.

**Exercise 4.13.** Describe  $\mathbb{Z} \rtimes C_2$  where  $C_2$  acts on  $\mathbb{Z}$  by inversion. How many sections are there of  $\mathbb{Z} \rtimes C_2 \rightarrow C_2$ ?

**Exercise 4.14.** How many sections are there to the projection  $D_{2n} \rightarrow C_2$ ?

## 5 AUG 30, 2023: $H^2$ , ABELIAN EXTENSIONS, AND BRAUER GROUP

Suppose we have an abelian extension, that is, let  $A$  be abelian, the short exact sequence of group extensions

$$0 \longrightarrow A \xrightarrow{i} E \xrightarrow{p} G \longrightarrow 1$$

is such that  $E/i(A) \cong G$ . Note that  $A$  can be regarded as a normal subgroup in  $E$  given this notation.

Note that two extensions are equivalent if there exists a group isomorphism  $\varphi : E \rightarrow E'$  such that the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \longrightarrow & E & \longrightarrow & G \longrightarrow 1 \\ & & \parallel & & \downarrow \varphi & & \parallel \\ 0 & \longrightarrow & A & \longrightarrow & E & \longrightarrow & G \longrightarrow 1 \end{array}$$

commutes.

Consider the continuous functions

$$\varphi : G \times G \rightarrow A$$

such that  $\varphi(g_1g_2, g_3) + \varphi(g_1, g_2) = \varphi(g_1, g_2g_3) + g_1\varphi(g_2, g_3)$ . We know  $H^2(G, M)$  is the quotient of all such functions over the coboundaries, i.e., the functions  $\varphi$  such that  $\varphi(g_1, g_2) = f(g_1) - f(g_1g_2) + g_1f(g_2)$ .

Now  $E \cong A \times G$  can be considered as a bijection, so we pick a set-theoretic section  $s : G \rightarrow E$  with  $s(1) = 1$ , and now every element in  $E$  is written as  $as(g)$  uniquely for some  $a \in A$  and  $g \in G$ , we have

$$s(g)a = s(g)as(g)^{-1}s(g) = {}^g as(g).$$

Note that  $s$  may not be a homomorphism, but we have  $s(g)s(h) = f(g, h)s(gh)$  since  $s(g)s(h)$  and  $s(gh)$  are both lifts of  $gh$ .

As a consequence, we have

$$(s(g_1)s(g_2))s(g_3) = f(g_1, g_2)s(g_1g_2)s(g_3) = f(g_1, g_2)f(g_1g_2, g_3)s(g_1g_2g_3)$$

and

$$s(g_1)(s(g_2)s(g_3)) = s(g_1)f(g_2, g_3)s(g_2, g_3) = {}^{g_1}f(g_2, g_3)s(g_1)s(g_2g_3) = {}^{g_1}f(g_2, g_3)f(g_1, g_2g_3)s(g_1g_2g_3).$$

In additive notation, we have

$$f(g_1, g_2) + f(g_1g_2, g_3) = g_1f(g_2, g_3) + f(g_1, g_2g_3).$$

Therefore,  $f$  becomes an inhomogeneous 2-cocycle.

**Proposition 5.1.** The induced map  $\lambda : \text{ext}(G, A) \rightarrow H^2(G, A)$  is a well-defined bijection between the set of equivalence classes of extensions and  $H^2(G, A)$ .



**Example 5.2.** The two elements in  $H^2(C_2, \mathbb{Z}/4\mathbb{Z}) \cong \mathbb{Z}/2\mathbb{Z}$  are given by non-split extension of  $Q_8$

$$0 \longrightarrow \mathbb{Z}/4\mathbb{Z} \longrightarrow Q_8 \longrightarrow C_2 \longrightarrow 1$$

and the identity element given by  $D_8 \cong \mathbb{Z}/4\mathbb{Z} \rtimes C_2$

$$0 \longrightarrow \mathbb{Z}/4\mathbb{Z} \longrightarrow D_8 \longrightarrow C_2 \longrightarrow 1$$

where  $D_8$  has the action of  $C_2$  over  $\mathbb{Z}/4\mathbb{Z}$ .

**Proposition 5.3.** An associative finite-dimensional  $K$ -algebra  $A$  is a CSA if and only if one of the following equivalent conditions hold:

1. Base-changed to the separable closure  $\bar{K}$  of  $K$  via  $\bar{K} \otimes_K A$ ,  $A \cong M_n(\bar{K})$  for some integer  $n \geq 1$ .
2. there exists a finite Galois extension  $L/K$  such that base-changed to  $L$  via  $L \otimes_K A$ ,  $A$  becomes isomorphic to a matrix algebra  $M_n(L)$  for some integer  $n \geq 1$ .
3.  $A \cong M_n(D)$  matrix algebra for some  $m \geq 1$  and some finite division algebra  $D$  over  $K$ .

A CSA  $A$  over  $K$  is said to be split over  $L$  if the above holds, i.e.,  $A \otimes_K L \cong M_n(L)$ . One can define an equivalence class on CSAs, such that  $A \sim B$  if and only if  $A \otimes_K M_n(K) \cong B \otimes_K M_m(K)$ . Now the Brauer group of  $K$  is the abelian group of equivalence classes of CSAs over  $K$  equipped with tensor product.

Suppose  $L/K$  is an extension, then there exists a homomorphism of base-change of algebras  $\text{Br}(K) \rightarrow \text{Br}(L)$ . We say the kernel  $\text{Br}(L | K)$  is the relative Brauer group of  $K$ -CSAs that split over  $L$ . The absolute Brauer group is  $\text{Br}(\bar{K} | K) = \text{Br}(K)$ , then

$$\text{Br}(K) = \bigcup_{L/K \text{ finite}} \text{Br}(L | K).$$

Now let  $L/K$  be a finite Galois extension with Galois group  $G$ , and we pick a normalized inhomogeneous 2-cycle  $\varphi : G \times G \rightarrow L^\times$  as the representative of its class, and we can construct  $A_\varphi$  as a  $K$ -CSA, then  $A_\varphi = \bigoplus_{g \in G} L e_g$  has dimension  $|G|^2$ , where  $e_g$ 's are the generators, with a multiplication operation  $(l e_g)(m e_h) = l(g \cdot m) \varphi(g, h) e_{gh}$  which can be extended via distribution.  $A_\varphi$  is said to be the crossed product of  $L$  and  $G$  via  $\varphi$ .

**Theorem 5.4.** 1.  $A_\varphi$  is a split algebra over  $L$ .

2. If  $\varphi, \varphi'$  are two normalized inhomogeneous 2-cocycles, then  $A_\varphi \sim A_{\varphi'}$  if and only if  $\varphi \sim \varphi'$ .
3.  $A_{\varphi\varphi'} \sim A_\varphi \otimes_K A_{\varphi'}$ .
4. Any  $K$ -CSA which is split over  $L$  is similar to a crossed product  $A_\varphi$  for some  $\varphi : G \times G \rightarrow L^\times$ .

**Corollary 5.5.**  $H^2(G, L^\times)$  is isomorphic to  $\text{Br}(L | K)$ , and  $H^2(\text{Gal}(\bar{K}/K), \bar{K}^\times)$  is isomorphic to  $\text{Br}(K)$ .

## 6 SEPT 1, 2023: COHOMOLOGY OF CYCLIC AND FREE GROUPS

Recall that we can compute  $H^*(G, M)$  using any acyclic resolution of  $M$ . We want to describe  $H^*(G, M)$  for specific  $G$  using nice resolutions.

We have

$$\dots \rightarrow G^3 \xrightarrow{\delta} G^2 \xrightarrow{\delta} G$$

and to obtain  $X^*(G, M)$  we map out of the resolution and into  $M$ , so  $\text{Map}(G, M) \cong \text{Hom}(\mathbb{Z}[G], M)$  as  $G$ -modules, and in general we obtain

$$\text{Map}(G^k, M) \cong \text{Hom}_{\mathbb{Z}}(\mathbb{Z}[G]^{\otimes k}, M)$$

as  $\mathbb{Z}$ -modules.

We denote  $F^{\text{st}}$  to be the standard free resolution given by

$$\mathbb{Z}[G]^{\otimes k} \xrightarrow{d} \mathbb{Z}[G]^{\otimes(k-1)} \rightarrow \dots \rightarrow \mathbb{Z}[G]^{\otimes 2} \xrightarrow{d_1-d_0} \mathbb{Z}[G]$$

To obtain  $X^*(G, M)$ , we can map this into  $M$ . Now the standard resolution becomes an augmentation of  $\mathbb{Z}$  that makes  $X^*(G, M)$  exact, free, and acyclic. The kernel of  $\mathbb{Z}[G] \rightarrow \mathbb{Z}$  is the augmentation ideal of  $G$  as of  $\mathbb{Z}[G]$ . Since this is a  $G$ -equivariant map, then the augmentation ideal is a  $G$ -submodule of  $\mathbb{Z}[G]$ , as a free abelian group generated by the set  $\{(g-1) \mid 1 \neq g \in G\}$ .

**Lemma 6.1.** If  $P_* \rightarrow \mathbb{Z}$  is any free resolution of  $\mathbb{Z}$  as a  $G$ -module, then for a  $G$ -module  $M$ , we have  $H^*(G, M) \cong H^*(\text{Hom}(P_*, M))^G$ .

*Proof.* Since each  $P_i$  is free, then  $\text{Hom}(P_i, M)$  is an acyclic module, so  $M \rightarrow \text{Hom}(P_*, M)$  is an acyclic resolution of  $M$ . Now apply Proposition 2.28 in the notes.  $\square$

**Remark 6.2.**  $H^*(G, M) \cong \text{Ext}_{\mathbb{Z}[G]}^*(\mathbb{Z}, M)$  as universal  $\delta$ -functors.

Now let  $C_n$  be the cyclic group of order  $n$ , generated by element  $g$ , then  $\mathbb{Z}[C_n] \cong \mathbb{Z}[g]/(g^n - 1)$ , so we have  $0 = g^n - 1 = (g-1)N_g$  in  $\mathbb{Z}[C_n]$  where  $N_g$  is the norm element  $N_g = 1 + g + \cdots + g^{n-1}$ , so we have a free resolution of  $\mathbb{Z}$ :

$$\cdots \longrightarrow \mathbb{Z}[C_n] \xrightarrow{1-g} \mathbb{Z}[C_n] \xrightarrow{N_g} \mathbb{Z}[C_n] \xrightarrow{1-g} \mathbb{Z}[C_n] \xrightarrow{\varepsilon} \mathbb{Z}$$

where augmentation  $\varepsilon$  sends  $g$  to 1. This allows us to compute the cohomology of any  $C_n$ -modules.

**Proposition 6.3.** Let  $M$  be an  $C_n$ -module, then

$$H^i(G, M) = \begin{cases} M^G, & i = 0 \\ \{m \in M \mid N_g m = 0\} / (1-g)M, & i > 0 \text{ odd} \\ M^G / N_g M, & i > 0 \text{ even} \end{cases}$$

*Proof.* Taking  $\text{Hom}(P_*, M)^G$  gives

$$\cdots \longleftarrow M \xleftarrow{1-g} M \xleftarrow{N_g} M \xleftarrow{1-g} M \longleftarrow \cdots$$

$\square$

**Remark 6.4.** If  $M$  has trivial action, then

$$H^i(G, M) = \begin{cases} M, & i = 0 \\ M[n], & i > 0 \text{ odd} \\ M/n, & i > 0 \text{ even} \end{cases}$$

where  $M[n]$  is the  $n$ -torsion in  $M$ .

Now if  $T = \mathbb{Z}$  be with generator  $t$ , then  $\mathbb{Z}[T]$  is isomorphic to the Laurent polynomials, so we have a resolution

$$0 \longrightarrow \mathbb{Z}[T] \xrightarrow{1-t} \mathbb{Z}[T] \longrightarrow \mathbb{Z}$$

since  $(1-t)$  is not a zero-divisor of  $\mathbb{Z}[T]$ . Therefore, taking  $\text{Hom}(P_*, M)^T$  gives

$$0 \longleftarrow M \xleftarrow{1-t} M$$

$$H^i(T, M) = \begin{cases} M^T, & i = 0 \\ M_T, & i = 1 \\ 0, & \text{otherwise} \end{cases}$$

Now let  $X$  be a set, and let  $G_X$  be the free group on  $X$ .

**Proposition 6.5.** The augmentation ideal  $I_X$  is a free  $\mathbb{Z}[G_X]$ -module, generated by the set  $\{(x-1) \mid x \in X\}$ , and so the exact sequence

$$0 \longrightarrow I_X \longrightarrow \mathbb{Z}[G_X] \longrightarrow \mathbb{Z} \longrightarrow 0$$

is a free resolution of  $\mathbb{Z}$  as a  $G_X$ -module.

*Proof.* As  $\mathbb{Z}$ -bases of  $I_X$ , we have  $\{(g-1) \mid g \in G_X\}$ , but  $\{h(x-1) \mid h \in G, x \in X\}$  is also a  $\mathbb{Z}$ -linear basis for  $I_X$ .  $\square$

**Remark 6.6.** Groups are free if and only if they have cohomological dimension 1.

## 7 SEPT 6, 2023: CUP PRODUCT

**Remark 7.1.** 1. A crossed homomorphism would be a group homomorphism when  $G$  has trivial action on  $M$ .

2. If  $X$  is an  $A$ -torsor, then there is a given  $G$ -action and a right  $A$ -action so that  $X \times A \rightarrow X$  is given by a diagonal action compatible to the  $G$ -action. Therefore,  ${}^g(x \cdot a) = {}^gx \cdot {}^ga$ .

**Definition 7.2.** Let  $A$  and  $B$  be  $G$ -modules, then there is a notion of tensor product  $A \otimes_G B$  as a  $G$ -module via the diagonal action  $g(a \otimes b) = ga \otimes gb$ . On the level of cochain, we have a cup product

$$\begin{aligned} C^p(G, A) \otimes C^q(G, B) &\xrightarrow{\sim} C^{p+q}(G, A \otimes B) \\ (\alpha : G^{p+1} \rightarrow A) \otimes (\beta : G^{q+1} \rightarrow B) &\mapsto (\alpha \smile \beta) \\ (g_0, \dots, g_{p+q}) &\mapsto \alpha(g_0, \dots, g_p) \otimes \beta(g_p, \dots, g_{p+q}) \end{aligned}$$

**Proposition 7.3.**  $\partial(\alpha \smile \beta) = (\partial\alpha) \cup \beta + (-1)^{|\alpha|} \alpha \smile \partial\beta$ .

**Corollary 7.4.** • If  $\alpha$  and  $\beta$  are cocycles, then  $\alpha \smile \beta$  is also a cocycle.

• If  $\alpha$  is a cocycle  $\beta$  is a coboundary, or vice versa, then  $\alpha \smile \beta$  is a coboundary. Indeed, if  $\beta = \partial\gamma$ , then  $\partial(\alpha \smile \gamma) = (-1)^{|\alpha|} \alpha \smile \beta$ .

Therefore, on the level of cohomology, we have a (bilinear) cup product as well:

$$H^p(G, A) \otimes H^q(G, B) \rightarrow H^{p+q}(G, A \otimes B)$$

**Example 7.5.** • If  $p = q = 0$ , then

$$\begin{aligned} H^0(G, A) \otimes H^0(G, B) &\cong A^G \otimes B^G \rightarrow H^0(G, A \otimes B) \cong (A \otimes B)^G \\ a \otimes b &\mapsto a \otimes b \end{aligned}$$

• By extending this property, we get a  $G$ -equivariant pairing  $A \otimes B \rightarrow C$  and therefore

$$H^p(G, A) \otimes H^q(G, B) \xrightarrow{\sim} H^{p+q}(G, C).$$

**Example 7.6.** Let  $R$  be a commutative ring, and if there is a  $G$ -action on  $R$ , then the multiplication  $m : R \otimes R \rightarrow R$  is  $G$ -equivariant, so we have a cup product

$$\smile : H^p(G, R) \otimes H^q(G, R) \rightarrow H^{p+q}(R)$$

This has the following properties:

1. This is natural in  $A, B$ , and  $C$ .
2. This is compatible with connecting homomorphism and exact sequences, that is,
  - Given short exact sequences

$$0 \longrightarrow A' \longrightarrow A \longrightarrow A'' \longrightarrow 0$$

and

$$0 \longrightarrow C' \longrightarrow C \longrightarrow C'' \longrightarrow 0$$

and pairing  $A \otimes B \rightarrow C$ , then this induces  $A \otimes B \rightarrow C'$  and in the quotients we have  $A'' \otimes B \rightarrow C''$ , so  $\delta(\alpha \smile \beta) = \delta\alpha \smile \beta$ , so we have a commutative diagram<sup>1</sup>

$$\begin{array}{ccccccc} A' \otimes B & \longrightarrow & A \otimes B & \longrightarrow & A'' \otimes B & \longrightarrow & 0 \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \end{array}$$

<sup>1</sup>This may require the assumption that the modules are flat.

and thus

$$\begin{array}{ccc} H^o(G, A'') \otimes H^q(G, B) & \longrightarrow & H^{p+q}(G, A'' \otimes B) \\ \downarrow \delta \otimes 1 & & \downarrow \delta \\ H^{p+1}(G, A') \otimes H^q(G, B) & \longrightarrow & H^{p+q+1}(G, A' \otimes B) \end{array}$$

• Given

$$0 \longrightarrow B' \longrightarrow B \longrightarrow B'' \longrightarrow 0$$

and

$$0 \longrightarrow C' \longrightarrow C \longrightarrow C'' \longrightarrow 0$$

and pairings

$$\begin{array}{ccccccc} A \otimes B' & \longrightarrow & A \otimes B & \longrightarrow & A \otimes B'' & & \\ \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & C' & \longrightarrow & C & \longrightarrow & C'' \longrightarrow 0 \end{array}$$

$$\text{so } \delta(\alpha \smile \beta) = (-1)^{|\alpha|} \alpha \smile \delta\beta.$$

*Proof.* Let  $\alpha = [a]$  for  $a : G^{p+1} \rightarrow A$  and  $\beta = [b]$  for  $b : G^{q+1} \rightarrow B''$ , then there is a lift  $\tilde{b} : G^{q+1} \rightarrow B \rightarrow B''$ . Then we have

$$\begin{array}{ccccccc} C^q/B^q(B') & \longrightarrow & C^q/B^q(B) & \longrightarrow & C^q/B^q(B'') & \longrightarrow & 0 \\ \downarrow \partial & & \downarrow \partial & & \downarrow \partial & & \\ 0 & \longrightarrow & Z^q(B') & \longrightarrow & Z^{q+1}(B) & \longrightarrow & Z^{q+1}(B'') \end{array}$$

and by the snake lemma we have a connecting homomorphism over group cohomologies.  $\square$

## 8 SEPT 8, 2023: RESTRICTION AND TRANSFER

Recall that we have a chain-level cup product, and we extend it to the level of cohomology. The cup product has the following properties:

1. If  $p = q = 0$ , then the cup product is the natural composition

$$A^G \otimes B^G \rightarrow (A \otimes B)^G \rightarrow C^G$$

2. Functoriality.

3. We have  $\delta(\alpha \smile \beta) = \delta(\alpha) \smile \beta$ , and incorporating this with the exact sequence, we have  $\delta(\alpha \smile \beta) = (-1)^{|\alpha|} \alpha \smile \delta(\beta)$ .

By the universal property of the tensor product, there exists a unique bilinear pairing that also satisfies these properties. To prove this, we use dimension-shifting.

**Remark 8.1.** Let  $M$  be a module, and map it into the induced module with an extended short exact sequence

$$0 \longrightarrow M \longrightarrow \text{Ind}^G(M) = \text{Map}(G, M) \cong \text{Hom}_{\mathbb{Z}}(\mathbb{Z}[G], M) \longrightarrow M_1 \longrightarrow 0$$

Taking the fixed points, we have

$$0 \longrightarrow M^G \longrightarrow (\text{Ind}^G(M))^G \longrightarrow (M_1)^G \longrightarrow H^1(G, M) \longrightarrow 0 \longrightarrow \dots$$

$$\dots \longrightarrow 0 \longrightarrow H^k(G, M_1) \xrightarrow{\cong} H^{k+1}(G, M)$$

Here  $(M_1)^G \rightarrow H^1(G, M)$  is a surjection. Now we know  $\delta : H^i(G, M_1) \rightarrow H^{i+1}(G, M)$  is a surjection for  $i = 0$ , and is an isomorphism for  $i > 0$ .

Proceeding inductively, we define

$$0 \longrightarrow M_i \longrightarrow \text{Ind}^G(M) \longrightarrow M_{i+1} \longrightarrow 0$$

If we start with  $A \otimes B \rightarrow C$ , then use property (3) repeatedly to the short exact sequence above, we get the uniqueness.

**Example 8.2.** Consider  $G = C_2$ , and consider the cohomology ring  $H^*(C_2, \mathbb{F}_2)$ . The action is obviously trivial. This induced the sequence with augmentation

$$0 \longrightarrow \mathbb{F}_2 \longrightarrow \mathbb{F}_2[C_2] \longrightarrow \mathbb{F}_2 \longrightarrow 0$$

The boundary map is  $\delta : H^i(C_2, \mathbb{F}_2) \rightarrow H^{i+1}(C_2, \mathbb{F}_2)$  is an isomorphism for all  $i$ .

We know  $H^i(C_2, \mathbb{F}_2) = \mathbb{F}_2\{x_i\}$ , so we can write  $x_{i+1} = \delta x_i$ . The product  $x_i \smile x_j = \delta^i x_0 \smile \delta^j x_0 = \delta^{i+j} x_0 \smile x_0 = \delta^{i+j} x_0 = x_{i+j}$ . Hence,  $H^*(C_2, \mathbb{F}_2) \cong \mathbb{F}_2[x]$  where  $x = |x_1|$ .

Note that

$$H^i(C_2, M) = \begin{cases} M^{C_2}, & i = 0 \\ \ker(N)/(\sim), & i \text{ odd} \\ M^{C_2}/N, & i > 0 \text{ even} \end{cases}$$

**Remark 8.3.** For odd prime  $p$ , we want to use the same method to calculate  $H^i(C_p, \mathbb{F}_p)$  with trivial action, then this is  $\{\mathbb{F}_p, i \geq 0\}$ . For instance, if we look at  $x_1 \smile x_1$ , then this is  $(-1)^{|x_1|} x_1 \smile x_1$ , so this gives  $2x_1 \smile x_1 = 0 \in H^2 = \mathbb{F}_p$ , so this gives  $x_1 \smile x_1 = 0$ . Note that  $H^*(C_p, \mathbb{F}_p) \cong \bigwedge(x_1) \otimes \mathbb{F}_p[y]$ .

We now talk about the functoriality in  $G$ . Given  $G_1$  acting on  $M_1$  and  $G_2$  acting on  $M_2$ , and say  $\varphi : G_1 \rightarrow G_2$  is a group homomorphism, and a map of modules  $f : M_2 \rightarrow M_1$ , then we say  $\varphi$  and  $f$  is a compatible pair of morphisms if for any  $g \in G_1$ , the diagram

$$\begin{array}{ccc} M_2 & \xrightarrow{f} & M_1 \\ \varphi(g) \downarrow & & \downarrow g \\ M_2 & \xrightarrow{f} & M_1 \end{array}$$

This gives a map  $C^*(G_2, M_2) \rightarrow C^*(G_1, M_1)$ , and hence a map on cohomology  $H^*(G_2, M_2) \rightarrow H^*(G_1, M_1)$ . For instance, if  $\varphi = \text{id}$ , we obtain the functoriality in  $M$ , as we previously saw. Similarly, if  $f = \text{id}$ , and  $M = M_2$  is a  $G_2$ -module, on which  $g_1 \cdot m = \varphi(g_1) \cdot m$ .

There are some special situations from the relations above.

1. Conjugation: let  $H \subseteq G$  be a subgroup, and we consider  $A$  to be a  $G$ -module, then there is restriction of  $G$ -action on  $A$  to  $H$ , so  $A$  becomes a  $H$ -module. Let  $B \subseteq A$  be a  $H$ -submodule in this sense. This is preserved by the action of  $A$ , but not necessarily by the action of  $G$ . For any  $g \in G$ , let the right conjugation be  $h^g = g^{-1}hg$  on  $h$ , and let  ${}^gH = gHg^{-1}$  on subgroup  $H$ . The compatible morphisms are now

$$\begin{aligned} {}^gH &\rightarrow H \\ h &\mapsto h^g \end{aligned}$$

and

$$\begin{aligned} B &\rightarrow gB \\ b &\mapsto gb \end{aligned}$$

Therefore, the induced maps on conjugation is given by  $(g)_* = H^*(H, B) \rightarrow H^*({}^gH, gB)$ . Therefore,  $(g_1g_2)_* = (g_1)_*(g_2)_*$ .

2. Inflation: suppose  $H \triangleleft G$  is a normal subgroup. We have the canonical map  $G \rightarrow G/H$ . Let  $A$  be a  $G$ -module, then  $G/H$  acts on  $A^H$ , and we look at the inclusion  $A^H \hookrightarrow A$ . Now  $\varphi : G \rightarrow G/H$  and  $f : A^H \hookrightarrow A$  are compatible, so on the level of cohomology, we get an inflation map

$$\inf_G^{G/H} : H^*(G/H, A^H) \rightarrow H^*(G, A).$$

If we look at  $H_1 \subseteq H_2 \triangleleft G$  where  $H_i \triangleleft G$ , we have  $G \rightarrow G/H_1 \rightarrow G/H_2 \cong (G/H_1)/(H_2/H_1)$ , then the inflation is

$$\inf_G^{G/H_1} \circ \inf_{G/H_1}^{G/H_2} = \inf_G^{G/H_2}.$$

3. Restriction: Let  $\varphi : H \hookrightarrow G$  and consider  $A$  as  $G$ -module and  $H$ -module respectively. There is now a restriction map

$$\text{res}_H^G : H^*(G, A) \rightarrow H^*(H, A)$$

Now if  $H_1 \subseteq H_2 \subseteq G$ , then

$$\text{res}_{H_1}^G = \text{res}_{H_1}^{H_2} \circ \text{res}_{H_2}^G$$

Inflation and restriction fit in a long exact sequence.

Finally, we discuss corestriction/transfer/norm. Let  $G$  be a finite group and let  $M$  be a  $G$ -module, then we have  $M^G \hookrightarrow M$  as inclusion. On the other way around, we have

$$\begin{aligned} \text{tr}/N : M &\rightarrow M^G \\ m &\mapsto \sum_{g \in G} gm. \end{aligned}$$

9 SEPT 11, 2023:

Let  $\varphi : G_1 \rightarrow G_2$  and  $f : M_2 \rightarrow M_1$  be compatible, then we denote  $(\varphi, f)^* = H^*(G_2, M_2) \rightarrow H^*(G_1, M_1)$ , with

$$G_1^{\times(*+1)} \longrightarrow G_2^{\times(*+1)} \longrightarrow M_2 \xrightarrow{f} M_1$$

such that it follows composition, and  $(\varphi, f)^*$  commutes with  $\delta$ , i.e.,

$$\begin{array}{ccccccc} 0 & \longrightarrow & M_2' & \longrightarrow & M_2 & \longrightarrow & M_2'' \longrightarrow 0 \\ & & \downarrow f & & \downarrow f & & \downarrow f \\ 0 & \longrightarrow & M_1' & \longrightarrow & M_1 & \longrightarrow & M_1'' \longrightarrow 0 \end{array}$$

and therefore we have a commutative square

$$\begin{array}{ccc} H^k(G, M_2'') & \xrightarrow{\delta} & H^{k+1}(G_2, M_2') \\ (\varphi, f)^* \downarrow & & \downarrow (\varphi, f)^* \\ H^k(G_1, M_1'') & \xrightarrow{\delta} & H^{k+1}(G, M_1') \end{array}$$

For  $\alpha \in C^k(M_2'')/B^k$ , we trace it back to  $\tilde{\alpha} \in C^k(M_2)/B_k$ , and  $\alpha$  is sent to  $Z^{k+1}(M_2'')$ , but now that means  $\tilde{\alpha}$  lands in the kernel of  $Z^{k+1}(M_2) \rightarrow Z^{k+1}(M_2')$ , so this is in  $Z^{k+1}(M_2')$ .

$$\begin{array}{ccccccc} C^k(M_2)/B_k & \longrightarrow & C^k(M_2'')/B_k & \longrightarrow & 0 \\ \partial \downarrow & & \downarrow \partial & & \\ 0 & \longrightarrow & Z^{k+1}(M_2') & \longrightarrow & Z^{k+1}(M_2) & \longrightarrow & Z^{k+1}(M_2'') \end{array}$$

Moreover, we have  $(\varphi, f)^*(\alpha \smile \beta) = (\varphi, f)^*\alpha \smile (\varphi, f)^*\beta$ , whenever the modules are compatible.

For transfer/corestriction, if  $H \subseteq G$  is a subgroup with finite index, and  $M$  is a  $G$ -module, then we have

$$\begin{aligned} \mathrm{tr}_G^H : M^H &\rightarrow M^G \\ m &\mapsto \sum_{g \in G/H} gm \end{aligned}$$

For instance, we have  $\mathrm{tr} : \mathbb{Z}^H = \mathbb{Z} \rightarrow \mathbb{Z}^G = \mathbb{Z}$  is multiplication by  $[G : H]$ . Note that  $H^*(X^*(G, M)^G) = H^*(G, M)$ , but  $H^*(X^*(G, M)^H) = H^*(H, M)$ , and the latter maps to the former cohomology structure via the transfer mapping. Hence, we have  $\mathrm{tr}_G^H : X^*(G, M)^H \rightarrow X^*(G, M)^G$  giving  $\mathrm{tr}_G^H \equiv \mathrm{cores}_G^H : H^*(H, M) \rightarrow H^*(G, M)$ . This is not a ring homomorphism.

**Remark 9.1** (Properties). 1.  $\mathrm{tr}$  commutes with  $\delta$ , that is, for a short exact sequence of  $G$ -modules (hence a short exact sequence of  $H$ -modules),

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

then we have

$$\begin{array}{ccc} H^k(H, C) & \xrightarrow{\delta} & H^{k+1}(H, A) \\ \mathrm{tr} \downarrow & & \downarrow \mathrm{tr} \\ H^k(G, C) & \xrightarrow{\delta} & H^{k+1}(G, A) \end{array}$$

2. If  $H_1 \subseteq H_2 \subseteq G$  are subgroups with finite indices, then  $\mathrm{tr}_G^{H_1} = \mathrm{tr}_G^{H_2} \mathrm{tr}_{H_2}^{H_1}$ .

3.  $\mathrm{tr}(\mathrm{res}(\alpha) \smile \beta) = \alpha \smile \mathrm{tr}(\beta)$ . Now given a pairing  $A \otimes B \rightarrow C$  of  $G$ -modules, with  $H \subseteq G$ , then

$$\begin{array}{ccccc} H^i(H, A) & \otimes & H^j(H, B) & \xrightarrow{\smile} & H^{i+j}(H, C) \\ \mathrm{res} \uparrow & & \downarrow \mathrm{tr} & & \downarrow \mathrm{tr} \\ H^i(G, A) & \otimes & H^j(G, B) & \xrightarrow{\smile} & H^{i+j}(G, C) \end{array}$$

*Proof Idea.* By dimension shifting, we reduce the case  $H^0$ , in which we have an explicit description. We have  $A^H \otimes B^H \rightarrow C^H$ , so for  $\alpha \in A^G$  and  $\beta \in B^H$ , we have  $\mathrm{tr}(\alpha \otimes \beta) = \sum_{g \in G/H} g(\alpha \otimes \beta) = \sum g\alpha \otimes g\beta = \alpha \otimes \sum_{g \in G/H} g\beta$ .  $\square$

**Example 9.2.** Let  $R$  be a commutative ring with a  $G$ -action, then the restriction  $\mathrm{res} : H^*(G, R) \rightarrow H^*(H, R)$  is a ring homomorphism, so  $H^*(H, R)$  is a  $H^*(G, R)$ -algebra. The opposite side has  $\mathrm{tr}$  is a map of  $H^*(G, R)$ -modules where the cohomology of  $H$  is given the module structure from the restriction. This induces the Frobenius reciprocity.

**Remark 9.3** (Other compatibilities). Let  $K \subseteq H \subseteq G$  be (normal) subgroups, then  $G \rightarrow G/K \rightarrow G/H$  are quotient maps. The restrictions of inclusions correspond to inflations of surjections: if  $K \triangleleft G$ , then  $G \rightarrow G/K$  and  $H \rightarrow H/K$ , so  $\mathrm{inf}_H^{H/K} \circ \mathrm{res}_{H/K}^{G/K} = \mathrm{res}_H^G \circ \mathrm{inf}_G^{G/K}$ . Note that the maps are contravariants. Moreover, we have  $\mathrm{inf}_G^{G/K} \circ \mathrm{cores}_{G/K}^{H/K} = \mathrm{cores}_G^H \circ \mathrm{inf}_H^{H/K}$ .

If  $H \triangleleft G$ , then  $\mathrm{res}_H^G \circ \mathrm{cor}_G^H = N_{G/H}$ ; also,  $\mathrm{cor}_G^H \circ \mathrm{res}_H^G = [G : H]$ .

## 10 SEPT 13, 2023: SPECTRAL SEQUENCE

Whenever  $G$  is not cyclic or  $Q_8$ , the group cohomology  $H^*(G, M)$  would not have a small resolution. We know there is a pullback diagram

$$\begin{array}{ccc} M & \longrightarrow & \prod_p M_p^n \\ \downarrow & & \downarrow \\ M_{\mathbb{Q}} & \longrightarrow & \prod_p (M_p^n)_{\mathbb{Q}} \end{array}$$

Here  $M_{\mathbb{Q}} = M \otimes_{\mathbb{Z}} \mathbb{Q}$  is the base-change, and  $M_p^n = \varprojlim_i M/p^i$  is the completion. For finite group  $G$ , we have  $H^*(G, M_{\mathbb{Q}}) = M_{\mathbb{Q}}^G$  if  $*$  = 0 and is trivial otherwise. Now we have the diagram

$$\begin{array}{ccc} H^*(G, M) & \xrightarrow{\text{res}} & H^*(\{e\}, M) \\ & \searrow |G| & \downarrow \text{tr} \\ & & H^*(G, M) \end{array}$$

where  $H^*(\{e\}, M)$  is  $M$  if  $*$  = 0 and is otherwise trivial. Note that if  $*$  > 0, then  $H^*(G, M)$  is annihilated by  $|G|$ . Let  $P \subseteq G$  be a Sylow  $p$ -subgroup, then if  $P$  is normal, then  $H^*(G, M_p^n) \cong H^*(P, M_p^n)^{G/P}$ . Therefore we have a normal series  $\cdots \triangleleft P_2 \triangleleft P_1 \triangleleft P$  with simple enough quotients, e.g., as abelian series. Therefore, we need ways to reassemble the cohomology.

For  $H \triangleleft G$  we know there is a  $G/H$ -action on  $H^*(H, M)$  via conjugation, so we can calculate  $H^*(G/H, H^*(H, M))$ , hence calculate  $H^*(G, M)$  using Lyndon-Hochschild-Serre spectral sequences.

We will first look at Bockstein spectral sequences. We start by looking at the sequence

$$\cdots \subseteq p^2\mathbb{Z} \subseteq p\mathbb{Z} \subseteq \mathbb{Z}$$

and factors each inclusion  $p^k\mathbb{Z} \subseteq p^{k-1}\mathbb{Z}$  via  $p^k(\mathbb{Z}/p\mathbb{Z})$ , then we have cohomology  $H^*(G, M/p)[p]$ , thus calculate  $H^*(G, M)$ . (Here the attachment by  $p$  is given by tensoring  $\mathbb{Z}[v_0]$  with grading  $p$ .) In general, we construct the abstract version as filtered cochain complex, with

$$\cdots \subseteq F^{p+1}C^* \subseteq F^pC^* \subseteq \cdots \subseteq C^*$$

so we can map each term to the graded version  $\text{gr}^p C^*$ . We denote the inclusions by  $i$  and the projections to the graded versions by  $\pi$ . The goal is to understand  $H^*(C^*)$  from the building blocks  $H^*(\text{gr}^* C^*)$ . There exists the factoring

$$\begin{array}{ccccccc} \cdots & \longrightarrow & H^q(F^{p+2}) & \xrightarrow{i} & H^q(F^{p+1}) & \xrightarrow{i} & H^q(F^p) \longrightarrow \cdots \\ & & \delta \uparrow & \swarrow \pi & \delta \uparrow & \swarrow \pi & \\ & & H^q(\text{gr}^{p+1}) & & H^q(\text{gr}^p) & & \end{array}$$

This is the  $E_1$ -page of the spectral sequence, given by  $E_1^{p,q} = H^q(\text{gr}^p)$ . We denote  $d_1 : H^q(\text{gr}^p) \rightarrow H^{q+1}(\text{gr}^{p+1})$  as the composition. Obviously  $d_1^2 = 0$ .

Now the  $E_2$ -page is given by  $H^*(E_1, d_1)$ . For  $a \in \ker(d_1)$ , the map  $i$  induces  $\tilde{\delta} \mapsto \delta a$  by lifting, so  $\pi(\tilde{\delta}a) \in H^{q+1}(\text{gr}^{p+2}) = E_1^{p+2, q+1}$ , with  $d_1(\pi(\tilde{\delta}a)) = \pi\delta\pi(\tilde{\delta}a) = 0$ . We then define  $d_2([a]) = [\pi(\tilde{\delta}a)] \in E_2$ . We then proceed inductively and find higher pages. This is usually done by calculating derived pages.

11 SEPT 15, 2023

Recall that: if  $H$  is a finite group,  $A$  is a finite  $H$ -module, then an extension of  $H$  by  $A$  is a group  $G$  such that

$$0 \longrightarrow A \longrightarrow G \longrightarrow H \longrightarrow 1$$

is exact, where the  $H$ -module structure on  $A$  is realized via conjugation  $h \cdot a = hah^{-1} \in G$ . We already know that the equivalence classes of extensions of  $H$  by  $A$  correspond to  $H^2(H, A)$ , where  $A \rtimes H$  corresponds to  $0 \in H^2(H, A)$ .

**Theorem 11.1.** Let  $p$  be an odd prime,  $|G| = p^{n+1}$ , and  $G$  contains  $\mathbb{Z}_q$  for  $q = p^n$  as a subgroup. If this is the case, then  $G$  is either  $\mathbb{Z}_{p^{n+1}}$ ,  $\mathbb{Z}_q \times \mathbb{Z}_p$ , or  $\mathbb{Z}_q \rtimes \mathbb{Z}_p$ , where the generator  $e \in H$  acts on  $1 \in \mathbb{Z}_q$  by  $e1e^{-1} = 1 + p^{n-1}$ . We denote  $H = \mathbb{Z}_p$  in this case.

*Proof.* We want to look at the short exact sequence

$$0 \longrightarrow \mathbb{Z}_q \longrightarrow G \longrightarrow H \longrightarrow 1$$

where  $H = \mathbb{Z}_p$ .



**Lemma 11.2.** If  $p$  is an odd prime, and there exists integer  $a$  such that  $a^p \equiv 1 \pmod{p^n}$  for  $n \geq 2$ , then  $a \equiv 1 \pmod{p^{n-1}}$ .

*Subproof.* This is trivial if  $a = 1$ . If  $a \neq 1$ , let  $d(a)$  be the largest possible integer  $d$  such that  $a \equiv 1 \pmod{p^d}$ . It suffices to show that  $d(a) \geq n - 1$ . By Fermat's Little theorem, we have  $d(a) \geq 1$ . We now want to show  $d(a^p) = d(a) + 1$ . Indeed, let  $a = 1 + p^d b$ , then using the binomial theorem, we have  $a^p = (1 + p^d b)^p = 1 + p^{d+1} b + \dots$  where the omitted terms have higher order of  $p^{d+2}$ . However,  $d(a^p) \geq n$ , so  $d(a) \geq n - 1$ . ■

Now let

$$0 \longrightarrow \mathbb{Z}_q \longrightarrow G \longrightarrow H \longrightarrow 1$$

be the extension with  $|H| = p$ , then the  $H$ -module of  $\mathbb{Z}_q$  is given by a map  $\varphi : H \rightarrow \text{Aut}(\mathbb{Z}_q) \cong \mathbb{Z}_q^\times$ . Since  $|H|$  is prime, then  $\varphi$  is either trivial or injective.

If  $\varphi$  is trivial, then  $h1h^{-1} = 1$  for all  $h \in H$ , so  $G$  is an abelian group. By the fundamental theorem of abelian groups, we know  $G$  is either  $\mathbb{Z}_{p^{n+1}}$  or  $\mathbb{Z}_q \times \mathbb{Z}_p$ .

If  $\varphi$  is injective, then  $n \geq 2$ , otherwise the size of  $H$  is larger than the size of the units. Given some element  $h \in H$  such that  $h1h^{-1} = k$ , then  $k^p \equiv 1 \pmod{p^n}$ . By Lemma 11.2,  $k = 1 + p^{n-1}b$  for some  $b \in \mathbb{Z}_p$ . Because  $\varphi$  is injective, then the image of  $\varphi$  has size  $p$ , but every element in the image has the form of  $k$ , therefore the image is just the set of such elements. Let  $e \in H$  be a generator such that  $e1e^{-1} = 1 + p^{n-1}$ . Now let  $A = \mathbb{Z}_q$  with this  $H$ -module structure, and it suffices to show that  $H^2(H, A) = 0$ , then we have the semidirect product only.

Since  $H$  and  $A$  are both cyclic groups, we write down the periodic resolution to be

$$A \xrightarrow{e-1} A \xrightarrow{N} A \xrightarrow{e-1} A \xrightarrow{N} A \longrightarrow \dots$$

where  $N$  is the norm element  $\sum_{h \in H} h$ . We know the action via  $e - 1$  on 1 is  $(e - 1) \cdot 1 = (1 + p^{n-1}) - 1 = p^{n-1}$ , so  $\ker(e - 1) = p\mathbb{Z}/q\mathbb{Z}$ ; the action via  $N$  is  $N \cdot 1 = \sum_{b \in \mathbb{Z}_p} (1 + p^{n-1}b) \equiv p \pmod{p^n}$ , therefore the image of the norm map is  $\text{im}(\mathbb{Z}) = p\mathbb{Z}/q\mathbb{Z}$  as well. Therefore,  $H^2(H, A) = 0$ . □

**Corollary 11.3.** If we have a  $p$ -group  $G$  with  $p \neq 2$ , then there is a unique subgroup of order  $p$  and a unique subgroup of index  $p$ .

Let  $H$  be a normal subgroup of  $G$ , then we consider the free  $\mathbb{Z}[H]$ -resolution

$$\mathbb{Z} \longleftarrow C_H^0 \longleftarrow C_H^1 \longleftarrow C_H^2 \longleftarrow \dots$$

and we can try turning it into a free  $G$ -resolution of  $\mathbb{Z}[G/H]$  by taking the tensor via

$$\mathbb{Z} \otimes \mathbb{Z}[G/H] \cong \mathbb{Z}/[G/H] \longleftarrow C_H^* \otimes \mathbb{Z}[G/H]$$

Because  $\mathbb{Z}[H] \otimes_{\mathbb{Z}[H]} \mathbb{Z}[G] \cong \mathbb{Z}[G]$ , then we have

$$\mathbb{Z}[G/H] \cong \mathbb{Z} \otimes_{\mathbb{Z}[H]} \mathbb{Z}[G] \longleftarrow C_H^* \otimes_{\mathbb{Z}[H]} \mathbb{Z}[G]$$

Now given an arbitrary free  $\mathbb{Z}[G/H]$ -resolution and we want to map the given resolution to it.

$$\begin{array}{ccccccc} \mathbb{Z} & \longleftarrow & D_{G/H}^0 & \cong & \mathbb{Z}[G/H] & \longleftarrow & D_{G/H}^1 & \cong & \mathbb{Z}[G/H]^m & \longleftarrow & \dots \\ & & \uparrow & & & & \uparrow & & & & \\ & & C_H^* \otimes_{\mathbb{Z}[H]} \mathbb{Z}[G] & \longleftarrow & \dots & \longleftarrow & (C_H^* \otimes_{\mathbb{Z}[H]} \mathbb{Z}[G])^m & & & & \end{array}$$

The vertical maps are resolved as  $G$ -modules by using the resolution of  $\mathbb{Z}[G/H]$ . We claim that there are horizontal maps that gives a double complex whose total complex is a resolution of  $\mathbb{Z}$  as a  $G$ -module.

**Example 11.4.** Consider the dihedral group  $D_{2n} \triangleright C_n$ , so  $D_{2n}/C_n \cong C_2$ . In particular, say  $D_{2n}$  is generated by  $\tau$  of order  $n$  and  $T$  of order 2, so  $C_n$  is generated by  $\tau$  and  $C_2$  is generated by  $T$ . Consider the resolutions

$$D^* : \mathbb{Z} \longleftarrow \mathbb{Z}[T]/(T^2 - 1) \xleftarrow{T-1} \mathbb{Z}[T]/(T^2 - 1) \xleftarrow{T+1} \mathbb{Z}[T]/(T^2 - 1) \xleftarrow{T-1} \mathbb{Z}[T]/(T^2 - 1) \xleftarrow{T+1} \cdots$$

and

$$C^* : \mathbb{Z} \longleftarrow \mathbb{Z}[\tau]/(\tau^n - 1) \xleftarrow{\tau-1} \mathbb{Z}[\tau]/(\tau^n - 1) \xleftarrow{N_\tau} \mathbb{Z}[\tau]/(\tau^n - 1) \xleftarrow{\tau-1} \cdots$$

and so on. Therefore we have an induced resolution given by

$$\mathbb{Z}[T]/T^2 \longleftarrow \mathbb{Z}[D_{2n}] \xleftarrow{\tau-1} \mathbb{Z}[D_{2n}] \xleftarrow{N_\tau} \mathbb{Z}[D_{2n}] \xleftarrow{\tau-1} \mathbb{Z}[D_{2n}] \xleftarrow{N_\tau} \cdots$$

Now let the sequence of  $D_{G/H}^n$ 's be of

$$\begin{array}{ccccccc} \mathbb{Z} & \longleftarrow & \mathbb{Z}[T]/T^2 & \xleftarrow{T-1} & \mathbb{Z}[T]/T^2 & \xleftarrow{T+1} & \mathbb{Z}[T]/T^2 & \xleftarrow{T-1} & \mathbb{Z}[T]/T^2 & \xleftarrow{T+1} & \cdots \\ & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \\ \cdots & \longleftarrow & \mathbb{Z}[D_{2n}] & \xleftarrow{T-1} & \mathbb{Z}[D_{2n}] & \xleftarrow{T+1} & \mathbb{Z}[D_{2n}] & \xleftarrow{T-1} & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & \tau-1 \uparrow & & \tau-1 \uparrow & & \tau-1 \uparrow & & \tau-1 \uparrow & & \\ \cdots & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & N_\tau \uparrow & & N_\tau \uparrow & & N_\tau \uparrow & & N_\tau \uparrow & & \\ \cdots & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & \uparrow & & \uparrow & & \uparrow & & \uparrow & & \\ \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \end{array}$$

The horizontal maps are hard to construct, they may look like  $\tau - 1$ , but we need to introduce signs at certain places.

12 SEPT 18, 2023

We will build the resolution out of this diagram, using double complexes, where horizontal differential  $\partial^v$  and vertical differential  $\partial^h$  satisfies  $\partial^v \partial^h + \partial^h \partial^v = 0$  between  $C^{i,j}$ 's. There now exists a total complex Tot with

$$(\text{Tot}^\oplus(C^{*,*}))_n = \bigoplus_{i+j=n} C^{i,j}$$

and

$$(\text{Tot}^\Pi(C^{*,*}))_n = \prod_{i+j=n} C^{i,j}$$

so each degree of the total complex is given by a collection of terms with the same fixed total degree. From the above, we have

$$\begin{array}{ccccccc} \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longleftarrow & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \xleftarrow{N_\tau} & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & \downarrow T+1 & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longleftarrow & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \xleftarrow{N_\tau} & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & \downarrow T-1 & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & \longleftarrow & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \xleftarrow{N_\tau} & \mathbb{Z}[D_{2n}] & \xleftarrow{\tau-1} & \mathbb{Z}[D_{2n}] & \longleftarrow & \cdots \\ & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\ \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \end{array}$$

One can fill in the diagram so that each square anticommutes, so that this becomes a double complex.

**Example 12.1.** If we calculate  $H^*(D_{2n}, \mathbb{F}_2)$ , we would find the differentials of the total complex to be zero, therefore the cohomology (after taking  $\text{Hom}(C^{*,*}, \mathbb{F}_2)$ ) is just determined by the number of copies in the total complex, enumerated on  $\mathbb{F}_2$ .

If we think of the quaternions  $Q_8$  instead, with the presentation  $\langle \tau, T \mid \tau^2 = T^2 = (\tau T)^2, \tau^4 = 1 \rangle$ , then we obtain

$$\begin{array}{ccccccccc}
 \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longleftarrow & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \xleftarrow{N_\tau} & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \longleftarrow & \cdots \\
 & & \downarrow T+1 & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longleftarrow & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \xleftarrow{N_\tau} & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \longleftarrow & \cdots \\
 & & \downarrow T-1 & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & \longleftarrow & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \xleftarrow{N_\tau} & \mathbb{Z}[Q_8] & \xleftarrow{\tau-1} & \mathbb{Z}[Q_8] & \longleftarrow & \cdots \\
 & & \downarrow & & \downarrow & & \downarrow & & \downarrow & & \\
 \cdots & & \cdots & & \cdots & & \cdots & & \cdots & & \cdots
 \end{array}$$

To make this a complex, we need to add notions of differentials, where we find a nullhomotopic map so that given a term in some degree and any term in the following degree, there exists a differential from the former to the latter.

13 SEPT 20, 2023

We think of  $H \triangleleft G$  with  $G \twoheadrightarrow G/H$ , then as we discussed before there are chains

$$\begin{array}{c}
 \mathbb{Z} \longleftarrow \mathbb{Z}[G/H] \longleftarrow \cdots \\
 \uparrow \\
 \mathbb{Z}[G] \\
 \uparrow \\
 \vdots
 \end{array}$$

and therefore this gives an anti-commute square

$$\begin{array}{ccc}
 C_{i,j} & \xleftarrow{\partial_h} & C_{i+1,j} \\
 \partial_v \uparrow & & \uparrow \partial_v \\
 C_{i,j+1} & \xleftarrow{\partial_h} & C_{i+1,j+1}
 \end{array}$$

where  $\partial_v$  and  $\partial_h$  are  $G$ -equivariant.

**Theorem 13.1.** In this situation, there are equivariant maps, where  $d_0 = \partial_v : C_{i,j} \rightarrow C_{i,j-1}$ ,  $d_2 : C_{i,j} \rightarrow C_{i-2,j+1}$ , and so on, with  $d_r : C_{i,j} \rightarrow C_{i-r,j+r-1}$ , so that these differentials commute with the augmentation maps  $\varepsilon_i : C_{i,0} \rightarrow B_i$ , that is,  $\varepsilon d_1^C = d_1^B \varepsilon$  and such that

$$\cdots \xrightarrow{\Sigma d_r} \bigoplus_{i+j=n} C_{i,j} \xrightarrow{\Sigma d_r} \bigoplus_{i+j=n-1} C_{i,j} \xrightarrow{\Sigma d_r} \cdots$$

is a free resolution of the trivial  $G$ -module  $\mathbb{Z}$ .

We will filter  $C_{*,*}$  by  $(F^p C_{*,*})_n = \bigoplus_{i+j=n, i \geq p} C_{i,j}$ , then  $\text{gr}^p = F^p / F^{p+1}$ , so the filtration (horizontally/vertically) gives a spectral sequence with page 2 as  $E_2^{p,q} = H^p(G/H, H^q(H, M))$ .

**Example 13.2.** Consider

$$0 \longrightarrow C_4 \longrightarrow Q_8 \longrightarrow C_2 \longrightarrow 0$$

with  $B_*$  given by  $\mathbb{Z}[C_2]$ 's, and  $C_{i,j} = \mathbb{Z}[Q_8]$ . The  $E_2$ -page is now  $H^p(C_2, H^q(C_4, \mathbb{Z}/2\mathbb{Z}))$ , and as  $\tau$  acts trivially on the resolution, then  $d_2 = \pm(\tau + 1)$  is the zero map on the spectral sequence. One can show that  $d_3 = \pm T$ . There will then be periodicity on the picture for  $d_4$  and so on.

Now the spectral sequence gives us  $H^p(G/H, H^q(H, M)) \Rightarrow H^{p+q}(G, M)$ , and therefore the  $E_\infty$ -page, with  $\text{gr}^* H^{p+q} \cong \bigoplus_{p+q} E_\infty^{p,q}$ . In the example above we see  $H^0(Q_8, \mathbb{Z}_2) \cong \mathbb{Z}_2$  since the filtration ends there;  $\text{gr}^* H^1(Q_8, \mathbb{Z}_2) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ ;  $\text{gr}^* H^2(Q_8, \mathbb{Z}_2) \cong \mathbb{Z}_2 \oplus \mathbb{Z}_2$ ;  $H^3 = \mathbb{Z}/2\mathbb{Z}$ . This describes a general picture of  $H^{4k+i}$ , and we can remove the graded version and yields the same result.

14 SEPT 22, 2023

We think of how  $H^p(G/H, H^q(H, M))$  turns into  $H^{p+q}(G, M)$ . We know  $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$ , and we consider total degree  $n$ .

- If  $n = 0$ , then  $H^0(G/H, H^0(H, M)) \cong H^0(G, M)$ .
- If  $n = 1$ , then we have a long exact sequence

$$\begin{array}{ccccccc} 0 \rightarrow H^1(G/H, H^0(H, M)) & \xrightarrow{\text{inf}} & H^1(G, M) & \xrightarrow{\text{res}} & H^0(G/H, H^1(H, M)) & \xrightarrow{d_2} & H^2(G/H, H^0(H, M)) \xrightarrow{\text{inf}} H^2(G, M) \rightarrow Q \rightarrow 0 \\ & & \downarrow & \nearrow & \downarrow & \nearrow & \\ & & \ker(d_2) & & \text{coker}(d_2) & & \end{array}$$

More generally, we get a filtration on  $H^n(G, M)$  with associated grading  $E_\infty^{p, n-p} \cong E_R^{p, n-p}$  for some  $R \gg 0$ . In the exact sequence above, we obtain

$$0 \longrightarrow H^1(G/H, H^0(H, M)) \cong E_\infty^{1,0} \xrightarrow{\text{inf}} H^1(G, M) \longrightarrow \ker(d_2) \cong E_\infty^{0,1} \longrightarrow 0$$

and correspondingly  $\text{coker}(d_2) = E_\infty^{2,0}$  with  $Q$  given by

$$\ker(d_2^{1,1}) \cong E_\infty^{1,1} \hookrightarrow Q \xrightarrow{\pi} \ker(d_3)^{0,2} \cong E_\infty^{0,2}$$

so that  $\text{res} = \pi\alpha$ . The edge maps are given by

$$\begin{array}{ccc} E_\infty^{n,0} & \hookrightarrow & H^n(G, M) \\ \uparrow & & \uparrow \text{inf} \\ E_2^{n,0} & = & H^n(G/H, H^0(H, M)) \end{array}$$

and

$$\begin{array}{ccc} H^n(G, M) & \twoheadrightarrow & E_\infty^{0,n} \\ & \searrow \text{res} & \downarrow \\ & & H^0(G/H, H^n(H, M)) \end{array}$$

**Example 14.1.** Consider giving  $H^p(C_2, H^q(C_2, \mathbb{Z}_2))$  to  $H^{p+q}(C_4, \mathbb{Z}_2)$ . The thing we want to calculate is the spectral sequence of

$$C^{p,q} = X^p(G/H, X^q(G, M)^{G/H}).$$

Given  $f_i \in C^{p_i, q_i}$ , we take

$$C^{p_1, q_1} \times C^{p_2, q_2} \xrightarrow{\sim} X^{p_1+p_2}(G/H, X^{q_1}(G, M)^H \otimes X^{q_2}(G, M)^H)^{G/H} \xrightarrow{\sim} X^{p_1+p_2}(G/H, X^{q_1+q_2}(G, M)^H)^{G/H}$$

and so  $d_r(x \smile y) = d_r(X) \smile y + (-1)^{|x|} x \smile d_r(y)$ . Therefore this satisfies some kind of Leibniz's rule. We conclude that  $E_2^{*,*} \cong \mathbb{F}_2[x, y]$ . Therefore the arrows takes on grid other than ones of the form  $x^{2n}$  and  $x^{2n}y$ , which is given by the  $E_3$ -page and beyond. We conclude that  $E_4 \cong E_\infty = \mathbb{F}_2[x^2] \otimes \bigwedge(y)$ .

15 SEPT 25, 2023

We will work over  $\mathbb{F}_2$ -coefficients today. We were trying to calculate the spectral sequence via

$$1 \longrightarrow C_2 \longrightarrow C_{2^n} \longrightarrow C_{2^{n-1}} \longrightarrow 0$$

Here  $H^*(C_2) = \mathbb{F}_2[x]$  where  $|x| = 1$ .

**Proposition 15.1.**  $H^*(C_{2^n}) \cong \mathbb{F}_2[x_n, y_n]/(x_n^2)$  for some  $x_n \in H^1$  and  $y_n \in H^2$  and  $n > 1$ .

On the  $E_2$ -page, we need to move  $(0, 1)$  to somewhere so that the total degree 1 would have only one piece of information, so we move  $(0, 1)$  to  $(2, 0)$ , and similarly  $(n, 1)$  to  $(n+2, 0)$ . In general,  $E_\infty^{*,*} \cong E_3^{*,*} \cong \mathbb{F}_2[x^2] \otimes \mathbb{F}_2[x_{n-1}]/x_{n-1}^2$ . We identify the column of  $p = 1$  to be  $x_{n-1}$  and column of  $p = 2$  to be  $y_{n-1}$  and we identify  $y_{n-1} = x_{n-1}^2$ . In general,  $[f] \in E_\infty^{p,q}$  is equivalent to  $F^p H^*(G)/F^{p+1} H^*(G)$ , and given also  $[f'] \in E_\infty^{p',q'}$  for, then  $[f][f'] \in E_\infty^{p+p',q+q'}$ , then  $[ff'] = [f][f']$  modulo  $F^{p+p'+1} H^*(G)$ .

The edge maps are

$$H^k(G/H) \cong H^k(C_{2^{n-1}}) \xrightarrow{\text{inf}} H^k(G) \cong H^k(C_2) \xrightarrow{\text{res}} H^k(H) \cong H^k(C_2)$$

where  $\text{inf}$  is an isomorphism for  $k = 0, 1$  and zero otherwise, and  $\text{res}$  is an isomorphism for even  $k$ , and is zero otherwise.

Note that if  $G = \varinjlim G_i$  for finite groups  $G_i$ 's, then  $H^*(G) \cong \text{colim}_{i, \text{inf}} H^*(G_i)$ .

**Corollary 15.2.**  $H^*(\mathbb{Z}_2; \mathbb{F}_2) \cong \mathbb{F}_2[x]/x^2$  for  $x \in H^1$ .

If we think of  $H^*(D_{2n})$ , then we already have  $C_{2^{n-1}} \rightarrow D_{2n} \rightarrow C_2$ , so  $H^p(C_2, H^q(C_{2^{n-1}})) \Rightarrow H^*(D_{2n})$  already collapses. For  $n = 1$ , we have  $C_2$ ; for  $n = 2$ , we have  $C_2 \times C_2$  and resolve the cohomology by Kunnetth isomorphism  $H^*(C_2 \times C_2) \cong \mathbb{F}_2[x, y]$  for  $x, y \in H^1$ . For  $n \geq 3$ ,  $E_2^{*,*} \cong H^*(C_2) \otimes H^*(C_{2^{n-1}}) \cong \mathbb{F}_2[e] \otimes \mathbb{F}_2[x]/x^2 \otimes \mathbb{F}_2[y]$ . Since higher pages vanishes, this is also  $E_\infty^{*,*}$ . Let  $\mathcal{X} = [x] \in H^1(D_{2n})$ , and  $\mathcal{Y} = [y]$  and  $\mathcal{E} = [e]$ , then  $\mathcal{X}^2 \in \mathbb{F}_2\{\mathcal{X}, \mathcal{E}^2\}$ . Eventually this would be hard to compute, so we would look at something different.

If we think of  $D_8 \cong \langle T, \tau \mid T^2 = 1 = \tau^4, T\tau T = \tau' \rangle$ , then we have  $C_2 \cong \langle \tau^2 \rangle \rightarrow D_8 \rightarrow C_2 \times C_2$ . Similarly,  $E_2 \cong \mathbb{F}_2[e] \otimes \mathbb{F}_2[x, y]$ , where  $e^i$ 's are on position  $(1, i+1)$  and  $d_2(e) = \alpha x^2 + \beta y^2 + \gamma xy$ , so we obtain maps of spectral sequences to our sequence  $C_2 \cong \langle \tau^2 \rangle \rightarrow D_8 \rightarrow C_2 \times C_2$ , including

$$C_2 \longrightarrow C_2 \times C_2 \longrightarrow C_2 = \langle \tau T \rangle$$

$$C_2 \cong \langle \tau^2 \rangle \longrightarrow C_4 \longrightarrow C_2 \cong \langle \tau \rangle$$

$$C_2 \longrightarrow C_2 \times C_2 \longrightarrow C_2 \cong \langle \tau \rangle$$

When we say a map of spectral sequences we mean  $f^* : E_r^{*,*} \rightarrow \tilde{E}_r^{*,*}$  by sending  $d_r(x)$  to  $d_r(f^*x)$ , as maps of differential graded algebras. From one of the sequence above, we obtain

$$H^*(C_2, H^*(C_2)) \Rightarrow H^*C_2 \times C_2$$

with  $d_2(e) = 0$ . Take our original sequence with  $H^*(C_2, H^*(C_2 \times C_2)) \Rightarrow H^*(D_8)$ , we send this to above by  $e \mapsto e$ ,  $x \mapsto x$ , and  $y \mapsto 0$ , then by naturality (as we compare with the sequence above), we note  $d_2(e) = \alpha x^2 + \beta y^2 + \gamma xy$  where  $\alpha = 0$ ; similarly we note  $\beta = 0$  by comparing with another sequence. Therefore  $d_2(e) = \gamma xy$ .

16 SEPT 27, 2023

The cohomology rings  $H^*(G, F)$  we referred to today are with respect to  $F = \mathbb{F}_p$  where  $p$  is a prime.

**Theorem 16.1** (Evans-Venkov Theorem). For any finite group  $G$ , the cohomology ring  $H^*(G; \mathbb{F}_p)$  is Noetherian.

*Proof.* Suppose we know this holds for  $p$ -groups, then for an arbitrary group  $G$ , take its Sylow  $p$ -subgroup  $P \subseteq G$ . The cohomology rings give a restriction  $\text{res} : H^*(G) \rightarrow H^*(P)$  where  $H^*(P)$  is Noetherian. By assumption, we know  $\text{tr} : H^*(P) \rightarrow H^*(G)$  is the backwards mapping, and that  $\text{tr} \circ \text{res} = [G : P]$ , therefore this is an isomorphism. The transfer is then surjective and the restriction is injective. Therefore,  $H^*(G)$  is the subring of a Noetherian ring, then  $H^*(G)$  is Noetherian, as the retraction  $\text{tr}$  is fully faithful. Alternatively, we can show that  $I_1 \subseteq I_2 \subseteq \dots \subseteq H^*(G)$  stabilizes: we note that

$$\text{res}(I_1) \cup H^*(P) \subseteq \text{res}(I_2) \cup H^*(P) \subseteq \dots \subseteq H^*(P)$$

stabilizes. Let  $x \in \text{res}(I_k) \cup H^*(P)$ , i.e.,  $x = \text{res}(a_k) \cup b$  for some choices of  $a_k$  and  $b$ . Taking the transfer, we have  $\text{tr}(x) = \text{tr}(\text{res}(a_k) \cup b) = a_k \cup \text{tr}(b)$ . The point being  $I_k$ 's and  $(\text{res}(I_k) \cup H^*(P))$  are now composites to be an isomorphism, therefore we identify them to be the same. In particular, if  $a_j \in I_k \setminus I_{k-1}$ , so taking the restriction we end up in  $\text{res}(I_{k-1}) \cup H^*(P)$ , then sending it back via trace multiplies it by a unit, so it should end up in  $I_{k-1}$  again.

We now need to show that  $H^*(P)$  is Noetherian for all finite  $p$ -groups  $P$ . By an induction on order of  $P$ , for  $H^*(C_p) = \wedge(e) \otimes \mathbb{F}_p[y]$ , and given a central extension  $C_p \triangleleft P \twoheadrightarrow \bar{P}$ , we need to show that the statement holds for  $P$  given it holds for  $\bar{P}$ . We consider the spectral sequence  $E_2^{i,j} : H^i(\bar{P}, H^j(C_p)) \Rightarrow H^{i+j}(P)$ , the  $\bar{P}$ -action on  $H^j(C_p)$  is trivial since every action of  $p$ -group on  $\mathbb{F}_p$  is always trivial, therefore the  $E_2$ -page decomposes as the tensor product of two cohomology rings, so  $E_2^{*,*} = H^*(\bar{P}) \otimes_{\mathbb{F}_p} H^*(C_p) = H^*(P)[e, y]/e^2$ .  $E_2^{*,*}$  is Noetherian as a tensor product of two Noetherian rings. One can show that

- by induction, we can show that  $E_r^{*,*}$  is Noetherian (the kernel of each  $d_r$  map will be finitely-generated over  $E_r^{*,0}$  as an algebra), and
- moreover, there is  $N \gg 0$  such that  $E_N^{*,*} \cong E_\infty^{*,*}$ .

It then allows us to conclude that  $E_\infty$  is Noetherian, hence  $H^*(P)$  is Noetherian as well.  $\square$

Suppose we have a short exact sequence

$$0 \longrightarrow A \longrightarrow B \longrightarrow C \longrightarrow 0$$

of  $G$ -modules, then we obtain  $H^k(G, C) \rightarrow H^{k+1}(G, A)$  as a connecting homomorphism.

**Example 16.2.** Consider

$$0 \longrightarrow \mathbb{Z}_p \longrightarrow \mathbb{Z}_{p^2} \longrightarrow \mathbb{Z}_p \longrightarrow 0$$

then we obtain Bockstein  $\beta : H^k(G, \mathbb{Z}/p) \rightarrow H^{k+1}(G, \mathbb{Z}_p)$ . So we have  $\beta : H^*(G, \mathbb{F}_p) \rightarrow H^{*+1}(G, \mathbb{F}_p)$ . This map is

- natural in  $G$ ;
- a derivation, i.e.,  $\beta(x \cup y) = \beta x \cup y + (-1)^{|x|} x \cup \beta y$ ;
- $\beta^2 = 0$ .

These are called the Steenrod operations, with  $P^0 = \text{id} : H^*(G) \rightarrow H^*(G)$ , and  $P^i : H^*(G) \rightarrow H^{*+2(p-1)i}(G)$ , satisfying

1. if  $|x| = 2k$ , then  $P^k(x) = x^p$ ,
2. if  $|x| < 2k$ , then  $P^k(x) = 0$ , and
3.  $P^k(x \cup y) = \sum_{i=0}^k (P^i x) \cup (P^{k-i} y)$ .

**Example 16.3.** For example,  $H^*(C_p) \cong \wedge(e) \otimes \mathbb{F}_p[y]$ , with  $\beta(e) = y$ ,  $\beta(y) = 0$ , and  $p^1(y) = y^p$ .

17 SEPT 29, 2023

Let  $p$  be odd, and all coefficients are over the field  $\mathbb{F}_p$ . The Steenrod operations  $P^i$  for  $i \geq 0$  is given by

$$P^i : H^m(-) \rightarrow H^{m+2(p-1)i}(-)$$

satisfying

1.  $P^2 = \text{id}$ ;
2. if  $|x| = 2n$ , then  $P^n x = x^p$ ;
3. if  $|x| < 2n$ , then  $P^n x = 0$ ;
4.  $P^n(x \smile y) = \sum_{i+j=n} P^i x \smile P^j y$ ,

as well as the algebraic relations, e.g.,  $P^1 P^1 = 2P^2$ , as Adem relations.

**Definition 17.1** (Steenrod Algebra). The Steenrod algebra is  $A^* = \mathbb{F}_p \langle \beta, P^i, i \geq 1 \rangle / \sim$ , where  $\sim$  is given by Adem relations.

**Definition 17.2** (Milnor's  $Q_i$ -operations). Denote  $Q_0 = \beta$ ,  $Q_i = [P^{p^{i-1}}, Q_{i-1}]$ , e.g.,  $Q_1 = [P^1, \beta] = P^1 \beta - \beta P^1$ ;  $Q_2 = [P^p, P^1 \beta - \beta P^1] = P^p P^1 \beta + \dots$ . The key fact is that  $Q_i(x \smile y) = (Q_i x) \smile y + (-1)^{|Q_i||x|} x \smile Q_{i-1} y$ .

**Example 17.3.**  $H^*(C_p)$  is the exterior algebra  $\bigwedge(x) \otimes \mathbb{F}_p[y]$  where  $|x| = 1$  and  $|y| = 2$ , with  $\beta x = y$ . Then  $Q_1 x = (P^1 \beta - \beta P^1)(x) = y^p$ ;  $P^p y^p = y^{p^2} = Q_2 x$ . In general,  $Q_i x = y^{p^i}$ .

**Definition 17.4** (Fiber Bundle, Principal Bundle). A fiber bundle is the diagram  $F \rightarrow E \xrightarrow{\pi} B$ , where  $B$  is the base space,  $E$  is the total space, and  $F$  is the fiber, such that for any  $b \in B$ , there exists a neighborhood  $U$  of  $b$  such that  $\pi^{-1}(U) \simeq U \times F$ , with certain compatibility.

A principal  $G$ -bundle is a fiber bundle with fiber  $G$ . In this case,  $E$  inherits a free  $G$ -action.

**Remark 17.5.** If  $G$  is a finite group, then this gives a finite covering.

For a nice enough group  $G$ , there is a classifying space  $BG$  characterized by the fact that if  $X$  is a CW complex, then homotopy classes of map from  $X$  to  $BG$ , denoted  $[X, BG]$ , correspond to the principal  $G$ -bundles over  $X$ , such that there is a universal principal  $G$ -bundle

$$\begin{array}{c} EG \\ \downarrow \\ BG \end{array}$$

where  $EG$  is contractible, with the universal property that given  $f : X \rightarrow BG$ , there is a pullback  $f^* EG$  with respect to these maps.

**Remark 17.6.** • If  $G$  is a finite group, then  $\pi_k(BG) = \begin{cases} G, k = 1 \\ 0, k \neq 1 \end{cases}$  and therefore  $BG = K(G, 1)$ .

• For a group  $A$  and integer  $n \geq 0$ ,  $K(A, n)$  is a space with

$$\pi_m(K(A, n)) = \begin{cases} A, m = n \\ 0, m \neq n \end{cases}$$

If  $n \geq 2$ ,  $A$  needs to be abelian for these structures to exist.

**Example 17.7.** 1.  $B(G \times H) = BG \times BH$ .

2. If  $G = H \rtimes K$ , then the classifying space  $BG$  is isomorphic to the fiber product  $BH \times_K EK = (BH \times EK)/\Delta$  with respect to the diagonal  $K$ -action  $\Delta$ .

3. Let  $H^n = \prod_n H$  be a product of  $n$  copies of  $H$ . Permuting these  $H$ 's gives an action  $\Sigma_n$  on  $H^n$ , then there is the wreath product  $H^n \rtimes \Sigma_n = H \wr \Sigma_n$ . The classifying space  $B(H \wr \Sigma_n) \simeq (BG)^n \times_{\Sigma_n} E\Sigma_n$ . More generally, for a space  $X$ , we can permute the copies and get a fiber bundle

$$\begin{array}{c} X^n \times_{\Sigma_n} E\Sigma_n \\ \downarrow \\ B\Sigma_n \end{array}$$

where  $F = X^n$ . This bundle has a section

$$\begin{aligned} s : B\Sigma_n &\rightarrow X^n \times_{\Sigma_n} E\Sigma_n \\ s_x(y) &= (x, \dots, x, \tilde{y}). \end{aligned}$$

**Definition 17.8** (Serre Spectral Sequence). Given a fiber bundle  $F \rightarrow E \rightarrow B$ , there is a spectral sequence given by  $H^i(B, H^j(F)) \Rightarrow H^{i+j}(E)$ .

**Example 17.9.** For  $H \triangleleft G$ , the sequence  $BH \rightarrow BG \rightarrow B(G/H)$  gives the Lyndon-Hochschild spectral sequences.

**Example 17.10.** Consider  $X^p \rightarrow X^p \times_{C_p} EC_p \rightarrow BC_p$ , it gives

$$H^i(BC_p, H^j(X^p)) \Rightarrow H^{i+j}(X^p \times_{C_p} EC_p).$$

We have

$$H^*(BC_p, H^*(X^p)) \Rightarrow H^*(X^p \times_{C_p} EC_p).$$

where  $H^*(X^p) \cong H^*(X)^{\otimes p}$ , which decomposes as a direct sum of free and trivial terms. Let  $C_p = \langle T \rangle / (T^p - 1)$ . The free terms are generated by the image of  $1 + T + \dots + T^{p-1}$ , and the trivial terms are of the form  $x \otimes \dots \otimes x$ , i.e., fixed by the permutation action on  $C_p$ .

18 OCT 2, 2023

Again, we work on cohomology with coefficients in  $\mathbb{F}_p$ .

Let  $\Sigma_n$  act on  $X^n$  for some space  $X$ . (Similarly, the action of  $C_n$  on  $X^n$  gives  $X^n \times_{C_n} EC_n$ ) The space  $X^n \times_{\Sigma_n} E\Sigma_n$  has a free contractible  $\Sigma_n$ -space as  $\Sigma_n$ -fiber  $X^n \times E\Sigma_n$ . For instance, define  $H2\Sigma_n = H^n \rtimes \Sigma_n$ , then  $B(H2\Sigma_n) = (BH)^n \times_{\Sigma_n} E\Sigma_n$ . We will show that the spectral sequence for these collapses at  $E_2$ -page. Note that given a fibration  $F \rightarrow E \rightarrow B$ , there is a spectral sequence  $H^i(F, H^j(B)) \Rightarrow H^{i+j}(E)$ , for instance take  $H \triangleleft G \rightarrow G/H$ , then we have a fibration  $BH \rightarrow BG \rightarrow B(G/H)$ . For instance, take the fibration  $X^n \rightarrow X^n \times_{\Sigma_n} E\Sigma_n \xrightarrow{\pi} B\Sigma_n$ . This gives a spectral sequence  $H^i(\Sigma_n, H^j(X)^{\otimes n}) \Rightarrow H^{i+j}(X^n \times_{\Sigma_n} E\Sigma_n)$ . Note that  $\pi$  has a section  $s(y) = (x, \dots, x, \tilde{y})$ . Looking at the edge homomorphisms  $\pi^* : H^i(B\Sigma_n) \rightarrow E_{\infty}^{i,0} \rightarrow H^i(X^n \times_{\Sigma_n} E\Sigma_n)$ , there is also a retraction hence  $d_r : E_r^{*,*} \rightarrow E_r^{i,0}$ 's are zero.

Let  $G$  be a finite group, then  $BG = K(G, 1)$ , so by definition  $\pi_n(BG)$  is  $G$  if  $n = 1$  and is zero otherwise. If  $A$  is abelian group, then there are (Eilenberg-MacLane) spaces  $K(A, n)$  for all  $n \geq 0$ , with  $\pi_k(K(A, n))$  being  $A$  if  $n = k$  and is zero otherwise.

**Remark 18.1.** • there is a fibration  $K(A, n-1) \rightarrow E \rightarrow K(A, n)$  where  $E$  is contractible. Therefore,  $K(A, n-1)$  is the loop space on  $K(A, n)$ .

- If  $X$  is a space and  $A$  is an abelian group, then  $H^n(X; A)$ , as a representable functor, is given by the homotopy classes  $[X, K(A, n)]$  of maps of spaces.
- $K(A, n)$  is an  $\infty$ -loop space.
- $\tilde{H}^m(\mathbb{F}_p, j)$  is 0 if  $m \leq j$ , is  $\mathbb{F}_p\{\iota_j\}$  if  $m = j$ .

Consider  $X^p \rightarrow X^p \times_{C_p} EC_p \rightarrow BC_p$ , so we have  $H^i(BC_p, H^j(X)^{\otimes p}) \Rightarrow H^*(X^p \times_{C_p} EC_p)$ .



**Lemma 18.2.** Let  $V$  be an  $\mathbb{F}_p$ -vector space, and let  $V^{\otimes p}$  be a space with cyclic permutation acting upon it, then  $V^{\otimes p}$  is isomorphic to a direct sum of free and trivial portions via action by  $C_p$ . The trivial portion is generated by the diagonal image  $(v \otimes \cdots \otimes v)$  for some  $v \in V$ ; the free portion is generated by the image of  $(1 + T + \cdots + T^{p-1}) = N_T$ , if we consider  $C_p = \langle T \rangle$ .

**Remark 18.3.**  $H^*(X)^{\otimes p} = \bigoplus_{j_1 + \cdots + j_p} H^{j_1}(X) \otimes H^{j_2}(X) \otimes \cdots \otimes H^{j_p}(X)$  and so  $H^*(C_p, V^{\otimes p}) = H^0(C_p, V^{\otimes p}) \oplus \cdots \oplus H^*(C_p, \text{diag})$ , where the first terms are image of norm maps, and the last term is the portion representing the fixed points.

**Exercise 18.4.** Show that classes in  $H^0(C_p, H^*(X^{\otimes p}))$  which are in the image of transfer are permanent cycles.

What about  $H^0(C_p, \mathbb{F}_p\{w \otimes \cdots \otimes w\}) \subseteq H^*(X)^{\otimes p}$ ? Let  $w \in H^j(X)$ , so  $w$  is represented by  $f_w : X \rightarrow K(\mathbb{F}_p, j)$ , so the pullback  $f_w^*(\iota_j) = w$ . We have a fiber diagram

$$\begin{array}{ccccc} X^p & \longrightarrow & X^p \times_{C_p} EC_p & \longrightarrow & BC_p \\ f_w^p \downarrow & & \downarrow & & \parallel \\ K(\mathbb{F}_p, j) & \longrightarrow & K(\mathbb{F}_p, j) \times_{C_p} EC_p & \longrightarrow & BC_p \end{array}$$

We interpret this as having the first few rows above the zeroth row as  $K(\mathbb{F}_p, j)$ , so all differentials vanishes in this class: in the reduced cohomology, we see the cohomology starts at  $m = j$ , everything below would be the image of transfer map, which gives as free summands and has no higher cohomology. Hence, the first non-zero differential would have been  $\iota_j^{\otimes p}$  onto the zeroth row, but this is not allowed since it has no higher cohomology, so when we pullback  $w$ , we have  $d_r(\iota_j^p) = 0$  and therefore  $d_r(w^{\otimes p}) = 0$ . By Leibniz rule, everything vanishes since this generates everything.

19 OCT 4, 2023

**Theorem 19.1** (Evans-Venkos).  $H^*(G, \mathbb{F}_p)$  is Noetherian if  $G$  is a finite group.

*Proof.* We reduce the proof to  $p$ -groups and induct on orders of  $G$ . This works for  $C_p$  as a base case. We can also extend  $C_p \triangleleft E \twoheadrightarrow G$  for some  $G$  with a smaller order than  $E$ , then there is a spectral sequence by  $H^i(G, H^j(C_p)) \Rightarrow H^{i+j}(E)$ . To run the induction, we need to know that

**Proposition 19.2.** The spectral sequence above collapses at a finite stage.

*Subproof.* Given  $C_p \triangleleft E \twoheadrightarrow G$ , we can write  $E = \prod_{i=1}^{|G|} g_i C_p$  for some  $g_i \in E$  as coset representatives of  $E/G$ . Note that this extension is central so the action on  $C_p$  is trivial, but not trivial on  $E$ . Now  $h \in G$  will permute the  $g_i C_p$ 's, so there is a group homomorphism  $G \rightarrow \Sigma_{|G|}$ , hence  $C_p^{|G|} \rtimes \Sigma_{|G|} = C_p \wr \Sigma_{|G|} \hookrightarrow E$ , and

$$\begin{array}{ccccc} C_p^{|G|} & \longrightarrow & C_p \wr \Sigma_{|G|} & \longrightarrow & \Sigma_{|G|} \\ \Delta \uparrow & & \uparrow & & \uparrow \\ C_p & \longrightarrow & E & \longrightarrow & G \end{array}$$

Therefore this gives a mapping of spectral sequences, from  $H^*(\Sigma_{|G|}, H^*(C_p^{|G|})) \Rightarrow H^*(C_p \wr \Sigma_{|G|})$  to  $H^*(G, H^*(C_p)) \Rightarrow H^*(E)$ . Now  $H^*(G)$  is  $\mathbb{F}_p[x]/(x^2) \otimes \mathbb{F}_p[y]$  where  $|x| = 1$  and  $|y| = 2$ . Therefore,  $H^*(G, H^*(G)) \cong H^*(G) \otimes \mathbb{F}_p[x, y]/(x^2)$ . Recall that the first spectral sequence collapses at  $E_2$ , and we want to see the second spectral sequence collapses at finite stage. Also note that  $H^*(G)$ , the bottom row of the spectral sequence, is all zeros, so we need to find the action on  $\mathbb{F}_p[x, y]/(x^2)$ . This corresponds to the zeroth column of the spectral sequence. Since  $y^{|G|} = f^*(y^{\otimes |G|})$ , then  $y^{|G|}$

is a permutation cycle in the spectral sequence  $H^*(G, H^*(C_p)) \Rightarrow H^*(E)$ . Hence,  $E_\infty^{*,*} \cong \mathbb{F}_p[y^{|G|}] \otimes \left( \bigoplus_{j < 2|G|} E_\infty^{i,j} \right)$ .

The rows are now  $y^{|G|}$ -cyclic, i.e.,  $1, x, y, xy, \dots, y^{|G|}$ , and arrows cannot cross this cycle anymore, since it is cyclic and would end up in the same class again. Therefore, the spectral sequence collapses at the  $2|G|$ -page. ■

□

**Definition 19.3.** An elementary abelian  $p$ -group is of the form  $C_p^{\times r}$ .

If  $G$  is a finite group, then we can approximate the spectral sequence over  $G$  by these elementary abelian  $p$ -groups.

**Theorem 19.4** (Quillen). If  $w \in H^*(G)$  is such that the restriction  $\text{res}(w) \in H^*(V)$  for all elementary abelian subgroup  $V$  of  $G$  is nilpotent, then  $w$  is nilpotent.

*Proof.* It suffices to show that if  $\text{res}(w) = 0 \in H^*(V)$  for all  $V$ , then  $w$  is nilpotent. This is because  $H^*(V) = \mathbb{F}_p[y_1, \dots, y_r] \otimes \wedge(x_1, \dots, x_r)$ , so any nilpotent element in  $H^*(V)$  squares to zero.

We can reduce this to the case where  $G$  is a  $p$ -group. If  $w \in H^*(G)$  is nilpotent, then the transfer  $\text{tr}(w) \in H^*(P)$  into Sylow  $p$ -subgroup is nilpotent, and vice versa (invertible).

We have an extension  $H \triangleleft G \rightarrow C_p$ , so we assume inductively we know the result for  $H$ . Take  $w \in H^*(G)$ , then  $\text{res}(w)$  to elementary abelian groups is nilpotent, so by the inductive procedure we know  $\text{res}(w) \in H^*(H)$  is nilpotent, then take  $w$  to some power and the restriction in  $H^*(H)$  would become zero. Therefore, we just need to show that if  $w \in \ker(\text{res}(H^*(G) \rightarrow H^*(H)))$ , then  $w$  is nilpotent.

If we regard  $H^*(H)$  of  $C_p$  as the zeroth column in the spectral sequence, then for  $w \in \ker(\text{res}_H^G)$ ,  $w \in F^1 H^*(G)$ , where  $F^i$  is the filtration on columns  $i$  and higher. □

20 OCT 6, 2023

Recall:

**Theorem 20.1.** Let  $G$  be a finite group, then if  $w \in H^*(G)$  is such that  $w$  restricts to a nilpotent element in the cohomology of elementary abelian subgroups of  $G$ , then  $w$  is nilpotent. That is,  $\text{res} : H^*(G) \rightarrow \varinjlim_{V \subseteq G} H^*(V)$  where  $V$ 's are elementary abelian, then kernel consists of nilpotent elements. That is,  $\text{res}$  is an  $f$ -isomorphism.

*Proof.* We reduced the proof to the case of  $p$ -groups, and we proceed inductively on  $H \hookrightarrow G \rightarrow C_p$ . If we consider the spectral sequence of  $H^*(C_p, H^j(H)) \Rightarrow H^{i+j}(G)$ , then the first row of the diagram would be  $1, x, y, xy, y^2, \dots$ , and note that every term starting from 2 has a factor of  $y$ .

Note that for any  $\Gamma$ -module  $M$ ,  $M$  an  $\mathbb{F}_p$ -vector space, then  $H^*(\Gamma, M)$  is a module over  $H^*(\Gamma, \mathbb{F}_p)$ , i.e.,  $M \otimes_{\mathbb{F}_p} \mathbb{F}_p \cong M$ , then  $H^*(C_p, H^i(H))$  is a module over  $H^*(C_p) \cong \wedge(x) \otimes \mathbb{F}_p[y]$ , then

**Claim 20.2.**  $E_2^{i \geq 2, *} = F^2(H^*(G)) \subseteq (y)$ .

We need to show that if  $w \in \ker(\text{res}(H^*(G) \rightarrow H^*(H)))$ , then  $w$  is nilpotent. The kernel of the restriction would be  $F^1(H^*(G))$ , so whenever  $w$  is in the kernel of the restriction,  $w^2 \in F^2 H^*(G)$ . Run an induction on  $r$  to show  $\smile [y] : E_r^{i,j} \rightarrow E_r^{i+2,y}$  is surjective for all  $i, j$ . This means some power of  $w$  will be divisible by the image of some class in  $H^1(G)$  over Bockstein  $\beta$ . Therefore, some power of  $w$  is divisible by all  $\beta(H^1(G))$ . (Note that  $H^1(G) = \text{Hom}(G, \mathbb{Z}/p\mathbb{Z})$  where  $G$  is a  $p$ -group, so this is non-trivial.) Therefore, this power of  $w$  is a product of  $(\beta x_i)$ 's. To see this, we note  $H_i \rightarrow G \xrightarrow{x_i} C_p$  has  $x_i$ 's as generators of  $H^1(G)$ . Let  $w \in H^*(G)$ , then we can assume inductively that some power of  $w$  restricts to 0 in every proper subgroup. From the spectral sequence for  $H_i \triangleleft G \xrightarrow{x_i} C_p$ , then this power of  $w$  is  $(\beta x_i) \cdots$ .

**Lemma 20.3.** Let  $G$  be a  $p$ -group. Then  $G$  is not elementary abelian if and only if there are non-zero classes  $v_1, \dots, v_k \in H^1(G)$  such that  $\beta(v_1)\beta(v_k) = 0$ .

*Subproof.* Consider  $G' = [G, G]G^p \rightarrow G \xrightarrow{x_1, \dots, x_r} C_p^{\times r}$  where  $x_1, \dots, x_r$  are generators of  $H^1(G)$ , and it suffices to check that the map  $G \rightarrow C_p^{\times r}$  is an  $H_1$ -isomorphism. Eventually, finding such  $v_i$ 's in  $H^1(G)$  is equivalent to having  $\beta(v_i)$  not linearly independent in  $H^2(G)$ . We have

$$H^1(C_p^{\times r}) \xrightarrow{\sim} H^1(G) \longrightarrow H^1(G') \xrightarrow{d_2} H^2(C_p^{\times r}) \longrightarrow H^2(G).$$

then the statement above is equivalent to  $d_2 \neq 0$ . This forces  $H^1(G)$  is zero, so we have an  $H^1$ -isomorphism as required. ■

Therefore, this power of  $w$  has to be zero. □

21 OCT 9, 2023

**Definition 21.1.** Let  $G$  be a finite group,  $M$  be a  $G$ -module. The norm map  $Nm_G : M \rightarrow M$  sends  $m$  to  $\sum_{g \in G} gm$ , so

$$\begin{array}{ccc} M & \xrightarrow{Nm_G} & M \\ \downarrow & & \uparrow \\ M_G & \xrightarrow{Nm_G} & M^G \end{array}$$

**Definition 21.2.**

$$\hat{H}^*(G, M) = \begin{cases} H_{-* - 1}(G, M), & * \leq -2 \\ \ker(Nm_G), & * = -1 \\ \text{coker}(Nm_G), & * = 0 \\ H^*(G, M), & * \geq 1 \end{cases}$$

**Example 21.3.** Let  $G = C_p$  and  $M = \mathbb{Z}$ , we have

$$\cdots \longrightarrow \mathbb{Z}[C_p] \xrightarrow{1-g} \mathbb{Z}[C_p] \xrightarrow{N_g} \mathbb{Z}[C_p] \xrightarrow{1-g} \mathbb{Z}[C_p] \xrightarrow{N_g} \mathbb{Z}[C_p] \xrightarrow{1-g} \mathbb{Z}[C_p] \xrightarrow{\varepsilon} \mathbb{Z}$$

where  $\varepsilon \cdot g \mapsto 1$ . We have

$$Nm_{C_p}(m) = \sum_{i=0}^{p-1} g^i m = \sum m = pm,$$

therefore  $\text{coker}(Nm) = \mathbb{Z}/p\mathbb{Z}$  and  $\ker(Nm) = 0$ . Therefore

$$\hat{H}^*(C_p, \mathbb{Z}) = \begin{cases} \mathbb{Z}/p\mathbb{Z}, & * \text{ even} \\ 0, & * \text{ odd} \end{cases}$$

More generally,

$$\hat{H}^*(C_p, M) = \begin{cases} M^G / N_g M, & * \text{ even} \\ \{m \in M : N_g M = 0\} / (1-g)M, & * \text{ odd} \end{cases}$$

**Definition 21.4.** A complete resolution  $F_*$  of  $G$  is an exact sequence

$$\cdots \longrightarrow F_1 \longrightarrow F_0 \xrightarrow{d_0} F_{-1} \longrightarrow \cdots$$

of finitely-generated free  $\mathbb{Z}[G]$ -modules along with an element  $e \in F_{-1}$  which is  $G$ -fixed and generates  $d_0$ .

To obtain a complete resolution, we get

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \xrightarrow{Nm_G} & \text{Hom}(F_0, \mathbb{Z}) \longrightarrow \cdots \\ & & & & \searrow \varepsilon & & \nearrow \varepsilon^* \\ & & & & & \mathbb{Z} & \end{array}$$

where  $e = \varepsilon^*(1)$ . Conversely, given a complete resolution  $F$ , because  $e$  is  $G$ -fixed,  $F_{-1}$  is  $\mathbb{Z}[G]$ -free,  $e$  generates a copy of  $\mathbb{Z} \subseteq F_{-1}$ . Therefore we have

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_0 & \xrightarrow{d_0} & F_{-1} & \longrightarrow & \cdots \\ & & \searrow \varepsilon & & \nearrow \mu & & \\ & & & & \mathbb{Z} & & \end{array}$$

for  $\varepsilon : F_+ \rightarrow \mathbb{Z}$  and  $\mu : \mathbb{Z} \rightarrow F$ .

**Definition 21.5.**  $\hat{H}^*(G, M) = H^*(\text{Hom}_G(\hat{F}_*, M))$ .

Intuitively, we can compare  $F^* \otimes_G M$ , so  $\text{Hom}(F, \mathbb{Z}) \otimes_G M \cong \text{Hom}_G(F, M)$ .

**Lemma 21.6.** Let  $F$  be a finitely-generated free  $\mathbb{Z}[G]$ -module, so  $Nm_{\mathbb{Z}[G]}(F \otimes M)_G \rightarrow (F \otimes M)^G$  is an isomorphism.

To connect this definition with the previous one, we consider  $\hat{F}_*$ ,  $\text{Hom}_G(\hat{F}_*, M)$  for  $n < 0$ , then  $\text{Hom}_G(F_n, M) \cong F^n \otimes M$ . We can write  $F^+$  as the complex  $F_* \rightarrow \mathbb{Z}$  with augmentation  $\varepsilon : F_0 \rightarrow \mathbb{Z}$ , and  $\text{Hom}((F^+)^{\times}, \mathbb{Z})$  as  $\mathbb{Z} \rightarrow F_{-1} \rightarrow F_{-2} \rightarrow \cdots$  where  $G_\mu : \mathbb{Z} \rightarrow F_{-1}$ . Therefore,  $\hat{H}^n = H_{-n-1}(G, M)$  for  $n \leq -2$  and is  $H^n(G, M)$  for  $n \geq -1$ .

**Lemma 21.7** (Shapiro).  $\hat{H}^*(H, M) \cong \hat{H}^*(G, \mathbb{Z}[G] \otimes_{\mathbb{Z}[H]} M)$  where  $H \subseteq G$  and  $M$  is an  $H$ -module.

For augmentation  $\varepsilon : P_* \rightarrow \mathbb{Z}$ , then let  $\tilde{P}_*$  be the cone of  $\varepsilon$ .

**Definition 21.8.** The Tate complex is  $T(G, M) = \tilde{P}_* \otimes \text{Hom}(P_*, M)$ .

In this sense, we can also define  $\hat{H}^*(G, M) = H_{-*}(T_*(G, M)^G)$ .

22 OCT 11, 2023

Let  $G$  be a finite group, a complete resolution would be

$$\begin{array}{ccccccc} \cdots & \longrightarrow & F_1 & \longrightarrow & F_0 & \longrightarrow & F_{-1} \longrightarrow \cdots \\ & & & & \searrow \varepsilon & & \nearrow \\ & & & & \mathbb{Z} & & \end{array}$$

so that  $\hat{H}^*(G, M) = H^*(\text{Hom}_G(F_*, M))$  and  $\hat{H}_*(G, M) = H_*(F_* \otimes_G M)$ . Observe that  $\hat{H}^*(G, \mathbb{Z}[G]) = 0$ . More generally, induced modules satisfy  $\hat{H}^*(G, \text{Ind}_G(M)) = 0$  and  $\hat{H}^*(G, \text{Ind}_G^H(M)) \cong \hat{H}^*(H, M)$ .

**Corollary 22.1** (Dimension Shifting). For any finitely-generated module  $M$ , there are  $K$  and  $Q$  with

$$0 \longrightarrow M \longrightarrow \text{Ind}_G(M) \longrightarrow Q \longrightarrow 0$$

and

$$0 \longrightarrow K \longrightarrow \text{Ind}_G(M) \longrightarrow M \longrightarrow 0$$

such that  $\hat{H}^i(G, M) \cong \hat{H}^{i+1}(G, K) \cong \hat{H}^{i-1}(G, Q)$ . (Recall that if  $M$  is a  $G$ -module, then  $\text{Ind}_G(U(M)) \cong_G \mathbb{Z}[G] \otimes M$ , where  $U$  is the forgetful functor and  $\mathbb{Z}[G] \otimes M$  has the diagonal action.

**Example 22.2.** Let  $G = C_n = \langle T \rangle$ , with  $y \in H^2(C_n, \mathbb{Z}) \cong \mathbb{Z}/n\mathbb{Z}$  be the generator. The exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}[C_n] & \xrightarrow{1-T} & \mathbb{Z}[C_n] \longrightarrow \mathbb{Z} \longrightarrow 0 \\ & & & & \searrow & & \nearrow \\ & & & & I & & \end{array}$$

where  $I$  is the augmentation ideal, as the kernel/cokernel of the sequences. Therefore  $\hat{H}^{i-2}(C_n, \mathbb{Z}) \cong \hat{H}^i(C_n, \mathbb{Z}) \cong \hat{H}^{i+2}(C_n, \mathbb{Z})$ .

Because the middle terms are free, this gives  $H^0(-, \mathbb{Z}) \rightarrow H^1(-, I) \xrightarrow{\cong} H^2(-, \mathbb{Z})$ .

**Theorem 22.3.** There is a unique product (i.e., for a pairing  $A \otimes B \rightarrow C$  of  $G$ -modules, we get a pairing  $\hat{H}^k(G, A) \otimes \hat{H}^m(G, B) \rightarrow \hat{H}^{k+m}(G, C)$ ) on  $\hat{H}^*$  satisfying

- on  $\hat{H}^0$ , it is induced by  $A^G \times B^G \rightarrow C^G$ , and that
- the connecting homomorphism  $\delta$  satisfies  $\delta(a \smile b) = \delta a \smile b + (-1)^{|a|} a \smile \delta b$ , and  $\delta(a \smile b) = (-1)^{|a||b|} \delta(b \smile a)$ .

*Proof.* Uniqueness is the direct result of dimension shifting. For existence, it suffices to construct a suitable pairing on standard Tate cochains. We build a standard resolution  $X_* \rightarrow \mathbb{Z}$  where  $X_i = \mathbb{Z}[G^{i+1}] \cong \mathbb{Z}[G]^{\otimes(i+1)}$  and so  $\hat{X}_*$  is the diagram given by

$$\begin{array}{ccc} X_* & \xrightarrow{\quad} & \text{Hom}(X_*, \mathbb{Z}) \\ & \searrow & \nearrow \\ & \mathbb{Z} & \end{array}$$

For  $i > 0$ ,  $X_{-i} \cong \mathbb{Z}[G]^{\otimes i}$ , so we need suitable maps  $\varphi_{p,q} : X_{p+q} \rightarrow X_p \otimes X_q$  for all  $p, q \in \mathbb{Z}$  because

$$\hat{C}^p(A) \otimes \hat{C}^q(B) = \text{Hom}_G(X_p, A) \otimes \text{Hom}_G(X_q, B) \rightarrow \text{Hom}_G(X_p \otimes X_q, C) \xrightarrow{\varphi_{p,q}^*} \text{Hom}_G(X_{p+q}, C) = \hat{C}^{p+q}(C).$$

This allows us to write down what  $\varphi_{p,q}$  is supposed to be. □

**Example 22.4.** Consider

$$\hat{H}^p(G, \mathbb{Z}) \otimes \hat{H}^{-p}(G, \mathbb{Z}) \rightarrow \hat{H}^0(G, \mathbb{Z})$$

given by  $f : G^{p+1} \rightarrow \mathbb{Z}$  and  $g : G^p \rightarrow \mathbb{Z}$  in  $\hat{H}^p(G, \mathbb{Z})$  and  $\hat{H}^{-p}(G, \mathbb{Z})$  respectively, then

$$(f \smile g)(\sigma_0) = \sum_{\tau_i \in G} f(\sigma_0, \dots, \sigma_p) \cdot g(\tau_p, \dots, \tau_1)$$

but actually

$$\hat{H}^p(G, \mathbb{Z}) \otimes \hat{H}^{-p}(G, \mathbb{Z}) \rightarrow \hat{H}^0(G, \mathbb{Z}) = \mathbb{Z}/|G|$$

is a perfect pairing, i.e.,  $\hat{H}^{-p}(G, \mathbb{Z}) \cong \text{Hom}(\hat{H}^p(G, \mathbb{Z}), \mathbb{Z}/|G|)$ .

**Remark 22.5.** Let  $R$  be a ring with a  $G$ -action, then  $H^*(G, R) \rightarrow \hat{H}^*(G, R)$  is a ring homomorphism.

For the case  $G = C_n$ , this gives  $H^*(G, \mathbb{Z}) \cong \mathbb{Z}[y]/ny \rightarrow \hat{H}^*(G, \mathbb{Z}) = \mathbb{Z}/n\mathbb{Z}[y^{\pm 1}]$ .

More generally, for any  $C_n$ -module  $M$ ,  $H^*(C_n, M) \rightarrow \hat{H}^*(C_n, M)$  is a map between a module over  $H^*(C_n, \mathbb{Z})$  and a module over  $\hat{H}^*(C_n, \mathbb{Z})$ . This map is therefore the inversion of  $y$  (due to the cup product structure). For instance,  $\hat{H}^*(C_p, \mathbb{Z}/p\mathbb{Z}) \cong (\mathbb{Z}/p\mathbb{Z}[x, y/x^2])[y^{-1}]$ .

For a general  $G$ , if we have an exact sequence

$$0 \longrightarrow \mathbb{Z} \longrightarrow F_{k-1} \longrightarrow \cdots \longrightarrow F_0 \longrightarrow \mathbb{Z} \longrightarrow 0$$

where  $F_i$ 's are  $G$ -free, then for  $y_k \in \hat{H}^k(G, \mathbb{Z})$ , then if we cup with  $y_k$ , we get an isomorphism  $\hat{H}^n(G, M) \cong \hat{H}^{n+k}(G, M)$ .

23 OCT 13, 2023

Recall that we have  $\hat{H}^i(G, \mathbb{Z}) \otimes \hat{H}^{-i}(G, \mathbb{Z}) \rightarrow \hat{H}^0(G, \mathbb{Z})$ . More generally,

**Proposition 23.1.** For a  $G$ -module  $M$ ,  $\hat{H}^i(G, M^\vee) \otimes \hat{H}^{-i-1}(G, M) \xrightarrow{\sim} \hat{H}^{-1}(G, \mathbb{Q}/\mathbb{Z})$  where we denote  $M^\vee = \text{Hom}(M, \mathbb{Q}/\mathbb{Z}) = \frac{1}{|G|}\mathbb{Z}/\mathbb{Z}$  is a perfect pairing.

*Proof.* Use dimension shifting to reduce it to  $i = 0$ , then check explicitly. Recall for cyclic group  $G$ , we have

$$\hat{H}^n(G, M) \otimes \hat{H}^2(G, \mathbb{Z}) \xrightarrow{\sim} \hat{H}^{n+2}(G, M)$$

from

$$0 \longrightarrow \mathbb{Z} \longrightarrow \mathbb{Z}[G] \xrightarrow{1-\sigma} \mathbb{Z}[G] \longrightarrow \mathbb{Z} \longrightarrow 0$$

(When regarding  $\mathbb{Z}[G]$ 's as free modules, we have the second cohomology by noting the coboundary occurs twice.) □

**Definition 23.2** (Class Module).  $C$  is called a class module if for all subgroups  $H$  of (finite group)  $G$ ,

1.  $H^1(H, C) = 0$ ;

2.  $H^2(H, C) = \mathbb{Z}/|H|$ , where the generator is called the fundamental class.

For any  $C$  and  $\gamma \in H^2(G, C)$ , i.e.,  $\gamma : G \times G \rightarrow C$  is an inhomogenous cocycle, we define  $C(\gamma) = C \oplus \bigoplus_{1 \neq g \in G} \mathbb{Z}b_g$  where  $b_g$  is a formal basis element. The  $G$ -action is given by  $g \cdot b_n = b_{gh} - g_g + \gamma(g, h)$  and  $b_1 = \gamma(1, 1)$ . The composition  $\gamma : G \times G \rightarrow C \rightarrow C(\gamma)$  is a coboundary. ( $\gamma = \delta\beta$ ,  $\beta(g) = b_g$ .) Therefore,  $\gamma \in \ker(H^2(G, C) \rightarrow H^2(G, C(\gamma)))$ . We have an exact sequence

$$\begin{array}{ccccccc} 0 & \longrightarrow & C & \longrightarrow & C(\gamma) & \longrightarrow & \mathbb{Z}[G] \longrightarrow \mathbb{Z} \longrightarrow 0 \\ & & & & \searrow & & \nearrow \\ & & & & I_G & & \\ & & \nearrow & & \searrow & & \\ & & 0 & & 0 & & \end{array}$$

$b_g \mapsto g-1$

which gives  $\hat{H}^0(G, \mathbb{Z}) = \mathbb{Z}/|G|\mathbb{Z} \xrightarrow{\cong} \hat{H}^1(G, I_G) \xrightarrow{\delta} \hat{H}^2(G, C)$ .

**Theorem 23.3.**  $\delta^2 : \hat{H}^n(H, \mathbb{Z}) \rightarrow \hat{H}^{n+2}(H, C)$  is  $\delta^2(x) = x \smile \gamma_H$ , where  $\gamma_H = \text{res}_H^G(\gamma)$ . Moreover, the following are equivalent:

1.  $C(\gamma)$  is cohomologically trivial.
2.  $C$  is a class module with fundamental class  $\gamma$ .
3.  $\delta^2$  is an isomorphism for all  $n$  and all  $H$ .

*Proof.* (1)  $\Rightarrow$  (2):  $\hat{H}^1(H, C) \cong \hat{H}^0(H, I_G) \cong \hat{H}^{-1}(H, \mathbb{Z}) = 0$  and  $\hat{H}^2(H, C) = \hat{H}^0(H, \mathbb{Z}) = \mathbb{Z}/|H|\mathbb{Z}$ .

(2)  $\Rightarrow$  (1): We have

$$0 = \hat{H}^1(H, C) \longrightarrow \hat{H}^1(H, C(\gamma)) \longrightarrow \hat{H}^1(H, I_G) \longrightarrow \hat{H}^2(H, C) \longrightarrow \hat{H}^2(H, C(\gamma)) \longrightarrow \hat{H}^2(H, I_G)$$

By dimension shifting on  $0 \rightarrow I_G \rightarrow \mathbb{Z}[G] \rightarrow \mathbb{Z}$ , we have  $\hat{H}^1(I_G) = \hat{H}^0(\mathbb{Z}) = \mathbb{Z}/|H|\mathbb{Z}$ , and so  $\hat{H}^2(H, C) = \mathbb{Z}/|H|\mathbb{Z}$ , but it follows by a zero map to  $\hat{H}^2(H, C(\gamma))$ , therefore the map  $\hat{H}^1(H, I_G) \rightarrow \hat{H}^2(H, C)$  is also the zero map. We then note that  $\hat{H}^1(H, C(\gamma)) = 0 = \hat{H}^2(H, C(\gamma))$ . This implies  $C(\gamma)$  is cohomologically trivial.  $\square$

**Theorem 23.4** (Nakayama-Tate). If  $C$  is a class module with fundamental class  $\gamma$ , then

$$\hat{H}^i(G, \text{Hom}(M, C)) \otimes \hat{H}^{2-i}(G, M) \xrightarrow{\sim} \hat{H}^2(G, C)$$

is a perfect pairing in the sense that  $\text{Hom}(\hat{H}^{2-i}(G, M), \mathbb{Q}/\mathbb{Z}) \cong \hat{H}^i(G, \text{Hom}(M, C))$ . Note  $\text{Hom}(\hat{H}^{2-i}(G, M), \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(\hat{H}^{2-i}(G, M, H^2(G, C)))$ .

24 OCT 16, 2023

For a class module  $C$ , choose the generator  $\gamma$  of  $\hat{H}^2(G, C)$ , so  $\gamma$  is represented by  $c : G \times G \rightarrow C$  and defines a map  $G^{\text{ab}} \rightarrow C^G/N_G C = \hat{H}^0(G, C)$ . Now we have  $\hat{H}^2(G, \mathbb{Z}) \otimes \hat{H}^0(G, C) \rightarrow \mathbb{Z}/|G|$ . Therefore, by connecting  $0 \rightarrow \mathbb{Z} \rightarrow \mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z} \rightarrow 0$ , we have  $\hat{H}^2(G, \mathbb{Z}) \cong \hat{H}^1(G, \mathbb{Q}/\mathbb{Z}) = (G^{\text{ab}})^\vee$ . Therefore  $G^{\text{ab}} = \hat{H}^2(G, \mathbb{Z})^\vee \cong \hat{H}^0(G, C)$ . Therefore,  $\gamma$  defines an isomorphism, with inverse extends to  $C^G \rightarrow G^{\text{ab}}$ .

**Remark 24.1.** If  $A$  is  $k$ -torsion, then  $\text{Hom}(A, \mathbb{Q}/\mathbb{Z}) \cong \text{Hom}(A, \mathbb{Z}/k\mathbb{Z})$ .

**Theorem 24.2.** Let  $G$  be a profinite group,  $G = \varprojlim_u G/uG$  where  $G/uG$  is finite, then  $H^*(G, M) \cong \text{colim}_u H^*(G/uG, M^u)$ .

By Tate cohomology,  $\hat{H}^{>0}(G, M) = H^{>0}(G, M)$  and for  $i \leq 0$  we have  $\hat{H}^i(G, M) = \varprojlim_{\text{deflation}} \hat{H}^i(G/u, M^u)$ .

Let  $P_* \rightarrow \mathbb{Z}$  be some projective/free  $G$ -resolution, so we obtain  $H_*((P_* \otimes M)/G) = H^*(\text{Hom}(P_*, M)^G) = H^*(G, M)$ .

For  $U \subseteq V \subseteq G$ , we have  $G/uG \twoheadrightarrow G/vG$ , then we define the deflation to be the composition of norm and coinflation,

$$\text{def} : H_j(G/uG, M^u) \cong H_j(P_* \otimes M^u)/(G/uG) \xrightarrow{\text{coinf}} H_j(G/vG, (M^u)/v) \xrightarrow{\text{norm}} H_j(G/v, M^v).$$

25 OCT 18, 2023

Let  $k$  be a number field, then we may study  $H^*(\text{Gal}(\bar{k}/k), -)$ . Over the localization  $k_p$ , we may want to study  $\text{Gal}(\bar{k}_p/k_p)$  in the same way as  $\mathbb{C}/\mathbb{R}$  with absolute Galois group  $C_2$ . Note that  $\text{Gal}(\bar{k}_p/k_p)$  has finite cohomological dimension. To do this, we have patched Tate cohomology by putting duality in  $\text{Gal}(\bar{k}_p/k_p)$  and periodicity for  $C_2$  together.

For finite groups, Tate cohomology gives  $H^*(G, \mathbb{F}_p) \rightarrow \lim_{V \subseteq G} H^*(V, \mathbb{F}_p)$ , where  $V$  is an elementary abelian subgroup, has nilpotent kernel and cokernel. This is based on  $H^*(C_p, \mathbb{F}_p) = \mathbb{F}_p[y] \otimes \bigwedge(x)$  and  $\hat{H}^*(C_p, \mathbb{F}_p) = \mathbb{F}_p[y^{\pm 1}] \otimes \bigwedge(x)$ . Another idea is that if  $\Gamma$  is any group, then we have  $H^*(\Gamma, \mathbb{F}_p) \rightarrow \lim_{G \subseteq \Gamma} H^*(G, \mathbb{F}_p)$  where  $G \subseteq \Gamma$  is a finite group. The question is how well does this approximate.

Farrell has the following version of Tate cohomology. We say  $\Gamma$  is of virtual cohomological dimension  $k$ , if there exists a finite index subgroup  $U \subseteq \Gamma$  with codimension  $k$ . If the virtual cohomological dimension of  $\Gamma$  is finite, then

1.  $\hat{H}^*(\Gamma, M) = H^*(\Gamma, M)$  for  $* > k$ ,
2. if the cohomological dimension of  $\Gamma$  is finite, then  $\hat{H}^*(\Gamma, M) = 0$ .

When  $G$  is finite, we have complete resolutions

$$\begin{array}{ccc} P_* & \xrightarrow{\quad} & \text{Hom}(P_*, \mathbb{Z}) \\ & \searrow & \nearrow \\ & \mathbb{Z} & \end{array}$$

of free  $\mathbb{Z}[G]$ -modules since  $\text{Hom}(\mathbb{Z}[G], \mathbb{Z}) \cong \mathbb{Z}[G]$ .

**Definition 25.1.** For any  $\Gamma$ , a complete resolution of  $\Gamma$  is an acyclic complex  $F_*$  of projective  $\Gamma$ -modules, as well as a projective resolution  $P_* \rightarrow \mathbb{Z}$  such that  $F_r \cong P_r$  for  $r \gg 0$ , then  $\hat{H}^*(\Gamma, M) = H^*(\text{Hom}_\Gamma(F_*, M))$ .

**Remark 25.2.** • There is a complete resolution such that  $F_n \cong P_n$  for all  $n$  greater than the virtual cohomological dimension of  $\Gamma$ .

- Any two complete resolutions are chain equivalent.

Note that if  $H^k(G, M) = 0$  for all  $k > n$ , then the cohomological dimension of  $G$  is  $n$ . This implies there is a projective resolution of  $\mathbb{Z} \leftarrow P_0 \leftarrow \cdots \leftarrow P_n \leftarrow 0$  and vice versa.

**Example 25.3.** If  $G$  has finite cohomological dimension,  $F_* = 0$ ,  $P_* \rightarrow \mathbb{Z}$  has finite projective resolution. This is a complete resolution.

**Lemma 25.4.** If  $G$  has finite cohomological dimension, then any acyclic complex  $F_*$  of projectives is chain contractible.

*Proof.* Take  $0 \rightarrow K \rightarrow F_k \rightarrow \cdots \rightarrow F_{k-n} \rightarrow B \rightarrow 0$ , then  $H^i(G, B) = H^{i+n}(G, K) = 0$ , so  $B$  is projective therefore  $B$  as the kernel of differentials, which indicates we have a splitting on the image of differentials. We have chain nullhomotopy.  $\square$

26 OCT 20, 2023

Recall that a complete resolution is  $(F_*, P_* \rightarrow \mathbb{Z})$  where  $F_*$  is an unbounded acyclic complex of projectives, and  $P_* \rightarrow \mathbb{Z}$  are projective resolutions. That means for  $G$  such that  $\text{vcd}(G) < \infty$ ,  $\hat{H}^*(G, M) = H^*(\text{Hom}_G(F_*, M))$   $F_* \cong P_*$  in high dimensions.

To construct this, let  $U \subseteq G$  be of finite cohomological dimension, say  $\text{cd}(U) = n = \text{vcd}(G)$ , take any  $P_* \rightarrow \mathbb{Z}$ , then this is projective as a  $U$ -resolution. Since the resolution has finite length, we can let  $K$  be the kernel of the final map and get an exact sequence of finite length

$$\cdots \longrightarrow K \longrightarrow P_{n-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow \mathbb{Z} \longrightarrow 0$$

In particular,  $K$  is  $U$ -projective. Therefore,

$$\cdots \longrightarrow P_{n+1} \longrightarrow P_n \longrightarrow K \longrightarrow 0$$

is a projective resolution of  $K$ .

Eventually we build  $K_1$  as the cokernel of  $K_0 \rightarrow \text{Map}(G/U, K)$ , and build  $K_i$  as the cokernel of  $K_{i-1} \rightarrow \text{coind}_U^G(K_{i-1})$  for  $i \geq 2$ .

**Remark 26.1.** Key features:

- $\hat{H}^*(G, M) \cong H^*(G, M)$  for  $* > \text{vcd}(G)$ , and
- $\hat{H}^*(G, M)$  can be computed from the cohomology of finite subgroups of  $G$ .

Properties:

- Long exact sequences
- Shapiro's lemma:  $\hat{H}^*(G, \text{Ind}_H^G M) = \hat{H}^*(H, M)$ .

To give a cup product structure, we need  $F_* \rightarrow F_* \hat{\otimes} F_*$  where  $(F_* \hat{\otimes} F_*)_n = \prod_{i+j=n} F_i \otimes F_j$ . It suffices to construct

$$\begin{array}{ccc} F_{2m} & \longrightarrow & F_m \otimes F_m \\ \downarrow & & \downarrow \\ P_{2m} & \longrightarrow & P_m \otimes P_m \end{array}$$

for  $m > \text{vcd}(G)$ . By manipulation, we get  $F_{2m} \rightarrow F_{m+k} \otimes F_{m-k}$  with dimension shifting.

27 OCT 23, 2023

Consider

$$0 \longrightarrow \mathbb{Z} \longrightarrow D_\infty \longrightarrow C_2 \longrightarrow 1$$

with non-trivial  $C_2$ -action on  $D_\infty$ . We claim that  $D_\infty$  and  $\mathbb{Z} \times C_2$  has isomorphic Farrell-Tate cohomology.

Let  $G = \mathbb{Z} \times C_2$ .

**Lemma 27.1.** If  $G_1$  has finite cohomological dimension, and  $G_2$  has finite virtual cohomological dimension, then  $P_* \otimes F_*$ , where  $P_*$  is a projective resolution of  $\mathbb{Z}$  as  $G_1$ -module, and  $F_*$  is a complete resolution of  $G_2$ , is a complete resolution of  $G_1 \times G_2$ .

**Corollary 27.2.**  $\hat{H}^*(G_1 \times G_2) \cong H^*(G_1) \otimes \hat{H}^*(G_2)$ .

**Example 27.3.**  $\hat{H}^*(\mathbb{Z} \times C_2, \mathbb{F}_2) = \mathbb{F}_2[e, x^{\pm 1}]/e^2$  where  $|e| = 1 = |x|$ .

For  $D_\infty$ , consider the spectral sequence  $\hat{H}^p(C_2, H^q(\mathbb{Z}, \mathbb{F}_2)) \Rightarrow \hat{H}^{p+q}(D_\infty)$ . Since  $H^q(\mathbb{Z}, \mathbb{F}_2) = \mathbb{F}_2[e]/e^2$ , then the only differential is  $d_2$ , so this collapses to  $\hat{H}^*(D_\infty)$ . The graded structure on this is  $\mathbb{F}_2[e, x^{\pm 1}]/e^2$ , with ring structure such that either  $[x][e] = [xe]$  or  $[x][e] = [xe] + [x^2]$ . (Turns out the second one is the multiplication structure.)

We now start talking about duality. Recall that  $H^*(G) \cong H^*(BG)$ , so if  $BG$  is an orientable compact manifold, then Poincare duality holds in  $H^*(G)$ .

**Example 27.4.** For  $G = \mathbb{Z}^{\oplus n}$ , we have  $BG = \prod_n S^1$ .

Let  $G$  be a group of finite cohomological dimension  $n$ , so there exists a projective resolution  $P_* \rightarrow \mathbb{Z}$  such that  $P_i = 0$  for  $i > n$ . Therefore  $H^n(G, -)$  is a right exact functor, so there exists  $M$  such that  $H^n(G, M) \neq 0$ . Take a free  $F \twoheadrightarrow M$  then  $H^n(G, F) \neq 0$ . Therefore,  $H^n(G, \mathbb{Z}[G]) \neq 0$ .

**Corollary 27.5.** The cohomological dimension of  $G$  is the maximal value  $n$  such that  $H^n(G, \mathbb{Z}[G]) \neq 0$ .

**Remark 27.6.**  $\mathbb{Z}[G]$  has a left and right  $G$ -action, so  $H^n(G, \mathbb{Z}[G])$  has a right  $G$ -action, hence we have a tensor product  $H^n(G, \mathbb{Z}[G]) \otimes_{\mathbb{Z}[G]} M$  for any (left)  $G$ -module  $M$ , with a map into  $H^n(G, M)$  by  $f \otimes m \mapsto (g \mapsto f(g)m)$ .



**Proposition 27.7.** If  $G$  has cohomological dimension  $n$ , and is of type FP, i.e., has a projective resolution  $P_* \rightarrow \mathbb{Z}$  with each  $P_i$  finitely-generated over  $\mathbb{Z}[G]$  and  $P_i = 0$  for all  $i > n$ , then  $H^n(G, \mathbb{Z}[G]) \otimes_{\mathbb{Z}[G]} M \rightarrow H^n(G, M)$  is an isomorphism for any  $M$ .

Let  $D = H^n(G, \mathbb{Z}[G])$ , then  $D \otimes M$  is a  $G$ -module via  $g \cdot (d \otimes m) = dg^{-1} \otimes gm$ , then  $D \otimes_{\mathbb{Z}[G]} M = (D \otimes M)_G = H_0(G, D \otimes M)$ .

*Proof.* As a natural transformation of right exact functors, this commutes with direct sums and general colimits, so it suffices to check for  $M = \mathbb{Z}[G]$ .  $\square$

This extends to an isomorphism

$$H_i(D \otimes M) \cong H^{n-i}(G, M).$$

If so,  $G$  is called a duality group.

**Theorem 27.8.** If  $G$  is FP with cohomological dimension  $n$ , then  $G$  is a duality group if and only if  $H^i(G, \mathbb{Z}[G]) = 0$  for  $i \neq n$  and  $H^n(G, \mathbb{Z}[G])$  is a torsion-free abelian group.

*Proof.* Suppose  $G$  is a duality group, then we have  $H_i(D \otimes M) \cong H^{n-i}(G, M)$ , so take  $M = \mathbb{Z}[G] \otimes \mathbb{Z}/k\mathbb{Z}$  of  $\mathbb{Z}[G]$ , then  $M$  is induced, hence  $D \otimes M$  is also induced. Therefore,  $H_{>0}(G, D \otimes M) = 0$ , so  $H^{\neq n}(G, M) = 0$ . Take

$$0 \longrightarrow \mathbb{Z}[G] \xrightarrow{k} \mathbb{Z}[G] \longrightarrow \mathbb{Z}[G] \otimes \mathbb{Z}/k\mathbb{Z} \longrightarrow 0$$

and therefore we have  $H^n(G, \mathbb{Z}[G]) \cong H^n(G, \mathbb{Z}[G])$  since  $H_1(G, D \otimes \mathbb{Z}/k\mathbb{Z}[G]) = H^{n-1}(G, \mathbb{Z}/k\mathbb{Z}[G]) = 0$ .

Now suppose  $T_i(M) = H^{n-i}(M)$ , then it is a homological  $\delta$ -functor, and  $T_{i>0}$  is effaceable, i.e., for all  $M$ , there exists  $F \rightarrow M$  such that  $T_i(F) = 0$ . Let  $U_i(M) = H_i(G, D \otimes M)$ , then this is also a homological  $\delta$ -functor that is effaceable for  $i > 0$ . By the previous theorem we know  $T_0 \cong U_0$ , we have the duality.  $\square$

28 OCT 25, 2023

Suppose  $G$  is a group with finite cohomological dimension  $n$ . Let  $D = H^n(G, \mathbb{Z}[G])$ , then  $H_0(G, D \otimes M) \cong H^n(G, M)$ .

We say  $G$  is a duality group if  $H_i(G, D \otimes M) \cong H^{n-i}(G, M)$ , which is equivalent to having  $H^*(G, \mathbb{Z}[G]) = \begin{cases} 0, * \neq n \\ D, * = 0 \end{cases}$

and  $D$  is torsion-free. (In particular, the Poincare duality is when  $D = \mathbb{Z}$ . In addition, we say it is an oreintable poicare duality group if  $D \cong \mathbb{Z}$  as  $G$ -modules.)

Now suppose  $G$  is virtual in addition with cohomological dimension  $N$ , i.e., there exists  $U \subseteq G$  such that  $[G : U] < \infty$  and has finite cohomological dimension.

We say  $G$  is a virtual duality group is there exists subgroup  $U \subseteq G$  of finite index such that  $U$  is a duality group. We have  $D_U = H^n(U, \mathbb{Z}[U]) \cong H^n(G, \mathbb{Z}[G])$ . (This holds as  $U$ -modules but has no information of  $G$ -action.) Therefore,  $G$  is a virtual duality group if and only if  $H^*(G, \mathbb{Z}[G])$  is 0 for  $* \neq n$  and is torsion-free for  $* = n$ .

**Example 28.1.**  $G = D_\infty$  is a virtual duality group with virtual cohomological dimension 1 and  $\mathbb{Z} \subseteq D_\infty$  is the infinite cyclic group as duality group with index 2.

**Example 28.2.** The classifying space  $B\mathbb{Z}$  of  $\mathbb{Z}$  is  $S^1$ , therefore  $\mathbb{Z}$  is a Poincare duality group. If  $G$  is a free group on  $k > 1$  generators, then  $BG$  is a wedge of  $k$  circles, thus  $H^0(G, \mathbb{Z}[G]) = 0$ ,  $H^1(G, \mathbb{Z}[G]) = D$ , and  $(D \otimes \mathbb{Z})_G \cong H^1(G, \mathbb{Z}) = \mathbb{Z}^k$ .

We say  $D$  is a dualizing module. Suppose  $G$  is a virtual duality group, what is the (co)homology of  $M$ ? We need to build a complete resolution for  $G$ . Take  $P_* \rightarrow \mathbb{Z}$  and  $Q_* \rightarrow D$  as projective resolutions. Note that  $H^*(\text{Hom}_G(P_*, \mathbb{Z}[G]))$  is  $D$  if  $* = n$  and is 0 otherwise. We will denote  $\bar{A} = \text{Hom}_G(A, \mathbb{Z}[G])$ . If we look at the complex

$$0 \rightarrow \bar{P}_0 \rightarrow \cdots \rightarrow \bar{P}_n \rightarrow \bar{P}_{n+1} \rightarrow \cdots$$

with  $\delta_n : \bar{P}_n \rightarrow \bar{P}_{n+1}$ , then there is an embedding  $\ker(\delta_n) \hookrightarrow \bar{P}_n$ , with  $\ker(\delta_n) \rightarrow D$ . Therefore, there is  $Q_0$  surjecting into  $D$ , therefore gives a lift into  $\ker(\delta_n)$ , thus this defines  $Q_0 \rightarrow \bar{P}_0$ . Using the acyclic complex, this gives lifts  $Q_i \rightarrow \bar{P}_{n-i}$  inductively as quasi-isomorphisms. Therefore, this gives an acyclic complex  $C_*$  of

$$\bar{P}_0 \oplus Q_{n-1} \rightarrow P_1 \oplus Q_{n-2} \rightarrow \cdots \rightarrow \bar{P}_{n-2} \oplus Q_1 \rightarrow \bar{P}_{n-1} \oplus Q_0 \rightarrow \bar{P}_n.$$

**Claim 28.3.**  $F_* = \bar{C}_*$  is a complete resolution for  $G$ .

*Proof.* This is given by

$$\cdots \rightarrow P_{n+1} \rightarrow P_n \rightarrow P_{n-1} \oplus \bar{Q}_0 \rightarrow \cdots \rightarrow P_0 \oplus \bar{Q}_{n-1} \rightarrow \bar{Q}_n \rightarrow \bar{Q}_{n+1} \rightarrow \cdots$$

□

**Corollary 28.4.** For  $m < -1$ , we have  $\hat{H}^m(M) \cong H_{n-m-1}(D \otimes M)$ . For  $n \geq m \geq -1$ , we have a long exact sequence by using image of transfer, as

$$\hat{H}^{-1}(M) \hookrightarrow H_n(D \otimes M) \rightarrow H^0(M) \rightarrow \hat{H}^0(M) \rightarrow \cdots \rightarrow H_0(D \otimes M) \rightarrow H^n(M) \rightarrow \hat{H}^n(M).$$

For  $m > n$ ,  $H^m(M) \cong \hat{H}^m(M)$ .

**Corollary 28.5.** If  $G$  is a duality group, then  $H^m(M) \cong H_{n-m}(D \otimes M)$ .

29 OCT 27, 2023

Let  $K$  be a non-Archimedean local field, as a finite extension over  $\mathbb{Q}_p$  or  $\mathbb{F}_p((t))$ . Suppose  $p \nmid n, m$ , then we have the intuition to denote  $\left(\frac{m}{p}\right) = 1$  if and only if  $x^n - m$  splits modulo  $p$ , which is equivalent to  $p$  splits in  $\mathbb{Q}(\sqrt[n]{m})/\mathbb{Q}$ , which is equivalent to  $\text{Frob}_{\mathbb{Q}(\sqrt[n]{m})/\mathbb{Q}}(p) = 1$ . Therefore, we want to define

$$\left(\frac{m}{p}\right) \sqrt[n]{m} = \text{Frob}_{\mathbb{Q}(\sqrt[n]{m})/\mathbb{Q}}(p) \sqrt[n]{m}$$

This gives a map

$$\begin{aligned} I_{\mathbb{Q}} &\rightarrow \text{Gal}(K/\mathbb{Q}) \\ p &\mapsto \text{Frob}_{K/\mathbb{Q}}(p) = \left(\frac{K/\mathbb{Q}}{p}\right) \end{aligned}$$

which factors over  $I_{\mathbb{Q}}/N_{K/\mathbb{Q}}(I_K)$ .

We want to prove that

**Theorem 29.1.** For any finite abelian extension  $L/K$ , we have an isomorphism

$$\varphi_{L/K} : K^\times / N_{L/K} L^\times \rightarrow \text{Gal}(L/K).$$

To do this, we will look at the commutative diagram

$$\begin{array}{ccc} K^\times & \xrightarrow{\varphi_K} & \text{Gal}(K^{\text{ab}}/K) \\ \downarrow & & \downarrow \\ H^0(\text{Gal}(L/K), L^\times) = K^\times / N_{L/K} L^\times & \xrightarrow{\varphi_{L/K}} & \text{Gal}(L/K) \end{array}$$

We will use the following notations:

- $\mathcal{O}_K$  as the ring of integers,
- $p_K = \pi_K \mathcal{O}_K$  with  $\pi_K$  being the uniformizer,
- $k = \mathcal{O}_K/p_K$ ,
- $U_K = \mathcal{O}_K^\times$ , and  $U_k^{(i)} = 1 + \pi_K^i \mathcal{O}_K$ . Therefore,  $U_k^{(i)}/U_k^{(i+1)} \cong k$ .

Therefore, we want

$$\varphi_K(\pi)|_{K^{un}} = \text{Frob}_K.$$

We will denote  $H^r(G, L^\times) =: H^r(L/K)$ . Suppose  $L/E/K$  is an intermediate extension, we have the inflation map

$$H^r(E/K) \rightarrow H^r(L/K).$$

Suppose  $L/K$  is unramified, then  $G \cong \text{Gal}(l/k)$ . By Hilbert Theorem 90,  $H^1(G, L^\times) = 0$  implies  $H^1(G, U_L) = 0$ . Therefore  $L^\times = \pi_L^\mathbb{Z} U_L \cong \mathbb{Z} \times U_L$ . We can start by calculating  $H^r(G, l^\times) = 0$  and  $H^r(G, l) = H^r(G, kG) = H^r(1, k) = 0$  by Shapiro's theorem. That means  $H^r(G, U_L) = 0$ . To see this, we look at the norm map

$$U_k^{(i)} / U_k^{(i+1)} \twoheadrightarrow U_L^{(i)} / U_L^{(i+1)}$$

For  $x \in U_k$ , there exists  $y_0 \in U_L$ , therefore  $xNy_0^{-1} \in U_k^{(1)}$ , and therefore there exists  $y_1 \in U_L^{(1)}$  such that  $x(Ny_0y_1)^{-1} \in U_k^{(2)}$ . Proceeding inductively,  $y = \prod y_i$  satisfies  $xNy^{-1} \in \bigcap U_k^{(i)}$ , and by completion this is just 1, so  $xNy^{-1} = 1$ . Hence,  $H^0(G, U_L) = 0$ . Recall that  $H^1(G, U_L) = 0$  as well, therefore (Tate) cohomology of  $U_L$  vanishes and we only care about  $\mathbb{Z}$  in  $L^\times = \mathbb{Z} \times U_L$ . This gives  $H^r(G, L^\times) \cong H^r(G, \mathbb{Z})$ , and therefore there is an invariant map

$$H^2(G, L^\times) \cong H^2(G, \mathbb{Z}) \cong H^1(G, \mathbb{Q}/\mathbb{Z}) \rightarrow \mathbb{Q}/\mathbb{Z}$$

defined by  $f \mapsto f(\alpha)$ . This means we have an isomorphism

$$\begin{array}{ccc} \text{Hom}(\text{Gal}(K^m/K) \cong \widehat{\mathbb{Z}}, \mathbb{Q}/\mathbb{Z}) & \xrightarrow{\cong} & \mathbb{Q}/\mathbb{Z} \\ \uparrow \hat{\mathbb{Z}} \rightarrow \mathbb{Z}/m\mathbb{Z} & & \uparrow \\ \text{Hom}(\text{Gal}(L/K) \cong \mathbb{Z}/m\mathbb{Z}, \mathbb{Q}/\mathbb{Z}) & \longrightarrow & \frac{1}{[L:K]} \mathbb{Z}/\mathbb{Z} \end{array}$$

Now suppose  $L/K$  is ramified, then  $H^2(\bar{K}/K) = H^2(K^{ur}/K)$  since  $\bar{K} \cong \text{Br}(\bar{K}/K) \cong \text{Br}(K)$  is the Brauer group, the group of central simple algebras under certain conditions. Let  $L$  be a finite extension of  $K$  in  $\bar{K}/L/K$ , then using the spectral sequence of

$$1 \longrightarrow \text{Gal}(L/K) \longrightarrow \text{Gal}(\bar{K}/K) \longrightarrow \text{Gal}(\bar{K}/L) \longrightarrow 0$$

we have

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^2(L/K) & \xrightarrow{\text{inf}} & H^2(\bar{K}/K) & \xrightarrow{\text{res}} & H^2(K/L) \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & * & \longrightarrow & H^2(K^m/K) & \longrightarrow & H^2(L^{ur}/L) \\ & & \downarrow \cong & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \frac{1}{[L:K]} \mathbb{Z}/\mathbb{Z} & \longrightarrow & \mathbb{Q}/\mathbb{Z} & \xrightarrow{[L:K]} & \mathbb{Q}/\mathbb{Z} \end{array}$$

As we denote  $G = \text{Gal}(L/K)$ , then we denote the subgroup  $H = \text{Gal}(L/E)$ . Therefore we have  $H^1(H, L^\times) = 0$  and  $H^2(H, L^2) \cong \mathbb{Z}/[L:E]\mathbb{Z}$  where  $L^\times$  is the class module. By Tate's theorem, we have an isomorphism

$$G^{\text{ab}} = H_1(G, \mathbb{Z}) = H^0(G, \mathbb{Z}) \xrightarrow{\cong} H^2(G, L^\times) = K^\times / N_L^\times$$

and we define  $\varphi_{L/K} : K^\times / N_L^\times \xrightarrow{\cong} \text{Gal}(L/K)^{\text{ab}}$  as its inverse. When  $L/K$  is finite abelian, then we have an isomorphism  $K^\times / N_L^\times \cong \text{Gal}(L/K)$ , so taking the colimit, we have

$$\begin{array}{ccc} K^\times & \xrightarrow{\varphi_K} & \text{Gal}(K^{\text{ab}}/K) \\ \downarrow & & \downarrow \\ K^\times / N_L^\times & \xrightarrow{\varphi_{L/K}} & \text{Gal}(L/K) \end{array}$$

where the bottom map is defined by  $\pi_K \mapsto \text{Frob}_{L/K}$  as the generator.

30 Nov 1, 2023

We want to show the following: let  $G$  be FP with finite virtual cohomological dimension, and suppose elementary abelian subgroups of  $G$  have rank at most 1, then  $\hat{H}^*(G, M) \cong \prod \hat{H}^*(N_G(V), M)$  with equivariant cohomology, where  $M$  is  $p$ -local, and the product runs through  $V$  as conjugacy classes of non-trivial elementary abelian subgroups. In particular, when  $*$   $>$   $\text{vcd}(G)$ , this is isomorphic to  $H^*(G, M)$ .

This is a consequence of a more general formula  $\hat{H}^*(G, M) \cong \hat{H}_G^*(|\mathcal{A}|)$ : let  $\mathcal{A}$  be a poset of non-trivial elementary abelian subgroups of  $G$ , with conjugation action, then  $|\mathcal{A}|$  is its geometric realization. In particular, the rank of VR is the number of generators.

Let  $X$  be a  $G$ -CW complex, intuitively,  $X$  has a cell decomposition which is respected by its  $G$ -action.

**Definition 30.1.** Let  $M$  be a  $G$ -module, then we define the equivariant cohomology by

$$H_G^*(X; M) = H^*(\text{Hom}_G(P_*, C^*(X; M)))$$

where  $P_*$  is a projective resolution of  $\mathbb{Z}$  and  $C^*(X; M)$  is a complex of abelian groups with a  $G$ -action.

**Example 30.2.** 1.  $X = *$  with trivial action, then  $H_G^*(*, M) = H^*(G, M)$ .

2.  $X = G/H$  with translation action,  $H_G^*(X; M) = H^*(H; M)$ .

To calculate this, we filter  $\text{Hom}_G(P_*, C^*(X; M))$  in two ways (over the double complex) and get two spectral sequences:

- $E_2^{p,q} = H^p(G, H^q(X; M)) \Rightarrow H_G^{p+q}(X; M)$ , and
- $E_1^{p,q} = \bigoplus H^q(G_\sigma, M) \Rightarrow H_G^{p+q}(X; M)$ , where the direct sum runs through orbits of  $p$ -cells in  $X$ , i.e., let  $G_\sigma$  be the stabilizer of a  $p$ -cell  $\sigma$ .

**Example 30.3.** If  $G$  acts on  $X$  freely, then  $H_G^*(X; M) = H^*(X/G; \tilde{M})$ ; where  $M$  has a  $G$ -action, so  $\tilde{M}$  is the local system over this action. In particular, if  $M$  has trivial  $G$ -action, then this is just  $M$ .

For Farrell-Tate cohomology, we can do something similar. Let  $F_*$  be a (Farrell-)Tate complete resolution for  $G$ , then  $\hat{H}_G^*(X; M) = H^*(\text{Hom}_G(F_*, C^*(X; M)))$ . We observe that if  $Y \hookrightarrow X$  is a  $G$ -subspace such that the isotropy group  $G_\sigma$  is trivial for every cell in  $X \setminus Y$ , then the inclusion generates an isomorphism  $\hat{H}_G^*(X, M) \cong \hat{H}_G^*(Y, M)$  by the spectral sequence.

31 Nov 3, 2023

Let  $X$  be a  $G$ -CW complex, let  $C^*X; M) = \text{Hom}(C_*X, M)$ , then  $H_G^*(X; M) = H^*(\text{Hom}_G(P_*, C^*(X; M)))$  where  $P_*$  are projectives. Similarly, we have  $\hat{H}_G^*(X; M) = H^*(\text{Hom}_G(F_*, C^*(X; M)))$  where  $F_*$  is a complete resolution. For orbits of  $p$ -cells  $\sigma$ ,  $C_p X = \bigoplus \mathbb{Z}[G/G_\sigma]$ , so the spectral sequence  $E_{1,\text{cell}}^{p,q} = H^q(G, C^p(X; M)) = \bigoplus H^q(G_\sigma, M)$ . This converges to the equivariant cohomology  $H_G^{p+q}(X; M)$  with filtrations in  $M$ .

**Proposition 31.1.** If  $Y \subseteq X$  is a  $G$ -subcomplex such that the cells in  $X \setminus Y$  are free (or have stabilizers of finite cohomological dimension), then  $\hat{H}_G^*(X; M) \cong \hat{H}_G^*(Y; M)$ .

*Proof.* Use the spectral sequence above, just take everything over equivariant  $\hat{H}$ . □

**Theorem 31.2** (Smith Theory). Let  $G = C_p$  and  $X$  be a finite-dimensional  $G$ -CW complex.

- (a) If  $H^*(X; \mathbb{F}_p)$  is finitely-generated, then so is  $H^*(X^{C_p}; \mathbb{F}_p)$ .
- (b) If  $H^*(X; \mathbb{F}_p) \cong H^*(*; \mathbb{F}_p)$ , i.e.,  $X$  is  $p$ -acyclic, then  $X^{C_p}$  is also  $p$ -acyclic.
- (c) If  $X$  is a homology sphere, i.e.,  $H^*(X; \mathbb{F}_p) \cong H^*(S^n; \mathbb{F}_p)$ , so is  $X^{C_p}$ .

*Proof.* Consider the inclusion  $X^{C_p} \hookrightarrow X$ , the fixed points are trivial, so by the proposition  $\hat{H}_{C_p}^*(X) \cong \hat{H}_{C_p}^*(X^{C_p})$ , but the latter has a trivial  $C_p$ -action, so as  $\text{Hom}_G(F_*, C^*(X; \mathbb{F}_p)) \cong \text{Hom}_G(F_*, \mathbb{Z}) \otimes_{\mathbb{Z}} C^*(X; \mathbb{F}_p)$ , therefore we have a Künneth isomorphism that makes  $\hat{H}_{C_p}^*(X^{C_p}) \cong \hat{H}^*(C_p; \mathbb{F}_p) \otimes_{\mathbb{F}_p} H^*(X^{C_p}; \mathbb{F}_p)$ . Consider the spectral sequence  $\hat{H}^s(C_p, H^q(X)) \Rightarrow \hat{H}_{C_p}^*(X)$ , then the differential contributing to the spectral sequence is given by  $\hat{H}^{s+q}$ . Therefore, if  $H^*(X)$  is finitely-generated, then  $\hat{H}_{C_p}^*(X) \cong \hat{H}^*(C_p) \otimes \hat{H}^*(X^{C_p})$  is finitely-generated, which forces  $\hat{H}^*(X^{C_p})$  to be finitely-generated. This proves (a). If  $X$  has the cohomology of a point, i.e.,  $H^*(X) = \mathbb{F}_p$ , the spectral sequence collapses to one line only, therefore the spectral sequence gives  $\hat{H}_{C_p}^*(X) = \hat{H}^*(C_p, \mathbb{F}_p) \cong \hat{H}^*(C_p) \otimes H^*(X^{C_p})$ .  $\square$

**Lemma 31.3.** Suppose  $Z$  is a  $G$ -CW complex, such that each stabilizer of a cell of  $Z$  is a non-trivial finite subgroup  $K$  of  $G$ , and the fixed points  $Z^K$  is acyclic, then  $Z$  is cohomologically equivalent (by zigzag) to the geometric realization  $|\mathcal{F}|$ , where  $\mathcal{F}$  is the poset of non-trivial finite subgroups of  $G$  with conjugation action.

**Example 31.4.** Let  $U \subseteq G$  be a finite index subgroup with finite cohomological dimension, then there is a finite-dimensional  $U$ -free contractible space  $EU$ . Form  $Y = \text{Map}_U(G, EU) \cong \prod_{G/U} EU$  to be

- contractible,
- finite-dimensional,
- stabilizer of any of its cells is finite,
- and  $Y^K \cong *$  for any finite  $K$ .

With this, let  $Y_0 = \bigcup_{K \in \mathcal{F}} Y^K$ .

32 NOV 6, 2023

**Lemma 32.1.** Let  $Z$  be a  $G$ -CW complex such that the stabilizer of each cell of  $Z$  is a non-trivial finite subgroup of  $G$ , and for each  $K \subseteq Z$  finite subgroup,  $Z^K \simeq *$ , then  $Z \simeq |\mathcal{F}(G)|$  equivariantly, the poset of non-trivial finite subgroups of  $G$ . In particular, if  $Z^K \simeq *$  is a cohomology isomorphism, then so is the isomorphism in our conclusion.

*Proof.* Note that  $Z = \bigcup_{K \in \mathcal{F}(G)} Z^K$  is a covering of  $Z$  by contractible subspaces,  $Z^{K_1} \cap Z^{K_2} = Z^{K_1 K_2} = \begin{cases} *, & \text{if } K_1 K_2 \in \mathcal{F}(G) \\ \emptyset, & \text{otherwise} \end{cases}$ .

We have a correspondence between  $Z$ , the Cech complex associated to this cover, as well as  $|\mathcal{F}(G)|$ .  $\square$

**Remark 32.2.** Suppose  $\text{vcd}(G) < \infty$ ,  $U \subseteq G$  has finite index, and  $\text{cd}(U) < \infty$ . Let  $Y = \text{Map}_U(G, EU)$  and  $\hat{H}_G^*(Y) \cong \hat{H}^*(G)$ . Let  $Z = \bigcup_{K \in \mathcal{F}(G)} Y^K$ , and  $Y \setminus Z$  has free action. Therefore  $\hat{H}^*(G) \cong \hat{H}_G^*(Y) \cong \hat{H}_G^*(Z) \cong \hat{H}_G^*(|\mathcal{F}(G)|)$ .

Observe that  $\hat{H}^*(G)_{(p)} \cong \hat{H}_G^*(|\mathcal{F}_p(G)|)_{(p)}$  where  $\mathcal{F}_p(G)$  is the set of non-trivial finite  $p$ -subgroups. Because we only need  $Z^K \simeq *$  in  $H^*(-)_{(p)}$ , we use restriction and transfer from  $p$ -Sylow.

**Theorem 32.3** (Quillen). The inclusion  $i : \mathcal{A}_p(G) \subseteq \mathcal{F}_p(G)$ , from poset of non-trivial elementary  $p$ -abelian subgroups of  $G$  to non-trivial finite  $p$ -subgroups, induces an  $G$ -equivalence  $|\mathcal{A}_p(G)| \simeq |\mathcal{F}_p(G)|$ .

This follows from

**Theorem 32.4** (Quillen's Theorem A). If  $X \rightarrow Y$  is a map of posets such that for each  $y \in Y$ ,  $X/y = \{x \in X \mid x \leq y\}$  or  $y \setminus X = \{x \in X \mid y \leq x\}$ , the slice category, is contractible, then  $|f| : |X| \rightarrow |Y|$  is an equivalence.

Let  $P \in \mathcal{F}_p(G)$ , then  $i/P = \mathcal{A}_p(P)$ . let  $B$  be simple  $p$ -torsion, i.e., maximal elementary abelian subgroup, of the center of  $p$ . As  $B$  is non-trivial, then

$$\begin{aligned} \mathcal{A}_p(P) &\rightarrow B \setminus \mathcal{A}_p(P) \\ A &\mapsto AB \end{aligned}$$

where slicing under  $B$  is given by  $C \in \mathcal{A}_p(P)$  such that  $B \in C$ .

33 Nov 8, 2023

**Theorem 33.1.** If  $G$  is a discrete group of FP type with finite virtual cohomological dimension  $G$ , then  $\hat{H}^*(G)_{(p)} \cong \hat{H}_G^*(|\mathcal{A}_p|)_{(p)}$ , where  $\mathcal{A}_p$  gives non-trivial elementary abelian  $p$ -subgroups of  $G$ .

**Claim 33.2.** Same is true for profinite groups.

**Example 33.3.** If the rank of elementary abelian  $V \subseteq G$  is at most 1, then

$$\hat{H}^*(G, M)_{(q)} \cong \prod_{\text{conjugacy classes of } V \cong C_p \hookrightarrow G} \hat{H}^*(N_G(V), M)_{(p)}.$$

This works for profinite groups as well. Reference for more result: [here](#).

**Definition 33.4** (Morava Stabilizer Group). For prime  $p$ , consider degree- $n$  extension  $\mathbb{Z}/p^n\mathbb{Z}$  of  $\mathbb{Z}/p\mathbb{Z}$ , then given by a multiplicative lift  $\mathbb{F}_p^\times \rightarrow \mathbb{Z}/p^n\mathbb{Z}^\times$  using Hensel's lemma, we have a commutative diagram

$$\begin{array}{ccc} \mathbb{Z}/p\mathbb{Z} & \longrightarrow & \mathbb{Z}/p^n\mathbb{Z} \cong \mathbb{Z}/p\mathbb{Z}(\xi_{p^n}) \\ \downarrow & & \downarrow \text{ (mod } p) \\ \mathbb{F}_p & \longrightarrow & \mathbb{F}_{p^n} \cong \mathbb{F}_p(\xi_{p^n}). \end{array}$$

Given a Frobenius map  $\sigma$  acting on  $\mathbb{F}_{p^n}$ , this lifts to a unique Frobenius acting on  $\mathbb{Z}/p^n\mathbb{Z}$ .

Consider a non-commuting polynomial ring  $\mathbb{Z}/p^n\mathbb{Z}\langle S \rangle$ , we get  $\mathcal{O}_n = \mathbb{Z}/p^n\mathbb{Z}\langle S \rangle / \langle S^n - p, Sa - a^\sigma S \rangle$  for  $a \in \mathbb{Z}/p^n\mathbb{Z}$ .

The  $n$ th Morava stabilizer group is  $\mathbb{S}_n = \mathcal{O}_n^\times$ , also known to be the  $\bar{\mathbb{F}}_p$ -automorphism group of the height- $n$  formal group laws over  $\mathbb{F}_p$ .

**Remark 33.5.** There is a valuation  $\nu$  on  $\mathcal{O}_n$  such that  $\nu(S) = \frac{1}{n}$  and  $\nu(p) = 1$ .

$\mathcal{O}_n$  is free of rank  $n$  over  $\mathbb{Z}/p^n\mathbb{Z}$ , and any element of  $x \in \mathcal{O}_n$  can be written as a sum  $x = x_0 + x_1S + \cdots + x_{n-1}S^{n-1}$  with  $x_i \in \mathbb{Z}/p^n\mathbb{Z}$  and  $x \in \mathcal{O}_n^\times$  if and only if  $x_0 \in \mathbb{Z}/p^n\mathbb{Z}^\times$ .

**Remark 33.6.** Considering  $\mathbb{S}_n$  as the automorphism group of formal group law  $\Gamma_n$ , then it acts over the universal deformations of  $\Gamma_n$ , which corresponds to  $(E_n)_* \cong \mathbb{Z}/p^n\mathbb{Z}[[u_1, \dots, u_{n-1}]] [u^{\pm 1}]$ , where  $u_i$ 's are of degree 0 and  $u$  is of degree  $-2$ .

We want to calculate  $H^*(\mathbb{S}_n, (E_n)_*)$ . Instead, we will try to compute  $\hat{H}^*(\mathbb{S}_n, (E_n)_*)$  for  $n = p - 1$ , with virtual cohomological dimension  $n^2$ , so this also will compute  $H^k(\mathbb{S}_n, (E_n)_*)$  for  $k > n^2$ . See Symonds' paper on Farrell-Tate.

We will verify elementary abelian  $p$ -subgroups have rank at most 1, then that means  $\hat{H}^*(\mathbb{S}_n, (E_n)_*) \cong \prod \hat{H}^*(N(C_p))$  as normalizers of  $C_p$ , and over  $n = p - 1$  they correspond to extensions of  $C_p$ .

34 Nov 10, 2023

The goal now is to identify  $\hat{H}^*(\mathbb{S}_n, (E_n)_*)$  as something computable by Symonds' work, where  $\mathbb{S}_n$  is the Morava stabilizer. Recall  $\mathbb{S}_n = \mathcal{O}_n^\times$  and  $\mathcal{O}_n \left[ \frac{1}{p} \right]$  is a division algebra over  $\mathbb{Q}_p$  with invariant  $\frac{1}{n}$  given by  $\mathcal{O}_n = \mathbb{Z}p^n\langle S \rangle / \langle S^n - p, Sa = a^\sigma S \rangle$ . Given the action on the commutative diagram mentioned last time, we define  $\mathbb{G}_n = \mathbb{S}_n \rtimes \text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ . In homotopy theory, the  $K(n)$ -local sphere spectrum  $S_{K(n)}^\circ$  extends to Lubin-Tate theory  $E_n$  with  $\mathbb{G}_n$  acting on it, so that the local sphere spectrum identifies as the homotopy fiber  $E_n^{h\mathbb{G}_n}$ . This induces  $H^*(\mathbb{G}_n, (E_n)_*) \Rightarrow \pi_* S_{K(n)}^\circ$ , and we will compute it by looking at  $H^*(\mathbb{S}_n, (E_n)_*)^{\text{Gal}}$ .

Therefore, we want to compute  $\hat{H}^*(\mathbb{S}_n, (E_n)_*)^{\text{Gal}}$ , which is just  $\hat{H}^*(N_{\mathbb{G}_n}(N), (E_n)_*)$ , where  $N$  is a finite subgroup.

**Example 34.1.** For  $n = 1$ ,  $\mathbb{S}_n = \mathbb{G}_n = \mathbb{Z}_p^\times$ . If  $p$  is odd, then  $\mathbb{Z}_p^\times \cong \mathbb{Z}/(p-1)\mathbb{Z} \times \mathbb{Z}_p$ , then  $\mathbb{Z}/(p-1)\mathbb{Z}$  has trivial Farrell-Tate cohomology since it is prime to  $p$ , and  $\mathbb{Z}_p$  has trivial Farrell-Tate cohomology because it has finite cohomological dimension. If  $p = 2$ , then  $\mathbb{Z}_2^\times = \mu_2 \times \mathbb{Z}_2$ , where  $\mu_2 = \{\pm 1\}$ , and  $\mathbb{Z}_2$  has zero Farrell-Tate cohomology because it has finite cohomological dimension. Now  $\hat{H}^*(\mathbb{Z}_2^\times, (E_1)_*) \cong \hat{H}^*(\mu_2, (E_1)_*)$ , where the left-hand side computes the

spectrum, and the right-hand side computes the real  $K$ -theory. Here  $(E_1)_* \cong \mathbb{Z}_p[\mu^{\pm 1}]$  for  $|u| = -2$ , and  $(E_1)_{2k} = \begin{cases} \mathbb{Z}_p, k \text{ even} \\ \mathbb{Z}_p(\text{sgn}), k \text{ odd} \end{cases}$ . We look at the extension

$$\mathbb{F}_{p^n}^\times \cong \mu_{p^n-1} \cong \langle \omega \rangle \hookrightarrow \mathbb{S}_n.$$

Take  $X = \omega^{\frac{p-1}{2}} S$ , then  $X^n = -p$  because  $X^2 = \omega^{\frac{p-1}{2}} S \omega^{\frac{p-1}{2}} S = (\omega^{\frac{p-1}{2}})^{\sigma} S^2$ , and proceeding inductively gives  $X^n = (\omega^{\frac{p-1}{2}})^{\sigma} (\omega^{\frac{p-1}{2}})^{\sigma^2} \cdots (\omega^{\frac{p-1}{2}})^{\sigma^{n-1}} S^n = -p$ , then  $\mathbb{Q}_p(X) \hookrightarrow \mathcal{O}_n \left[ \frac{1}{p} \right]$  assuming  $n = p-1$ , and we get  $\xi_p \in \mathcal{O}_n$ , so we identify  $\mathbb{Q}_p(X) \cong \mathbb{Q}_p(\xi_p)$ .

- If  $(p-1) \nmid n$ , then there is no finite  $p$ -torsion in  $\mathbb{S}_n$ .
- If  $n = p-1$ , then any finite  $p$ -subgroup of  $\mathbb{S}_n$  is isomorphic to  $C_p$ .
- For general  $n$ , the finite  $p$ -subgroups are  $C_{p^k}$  and/or  $Q_8$  if  $p = 2$ .

**Corollary 34.2.**  $\hat{H}^*(\mathbb{S}_n, -)$  is isomorphic to the product of  $\hat{H}^*(N_{\mathbb{S}_n}(V), -)$  where  $N$  is the normalizer, and the product runs over  $V \subseteq \mathbb{S}_n$  as conjugacy classes of elementary abelian subgroups.

**Example 34.3.** In case of  $n = p-1$ , we have  $C_p \subseteq \mathbb{S}_n \cong \mu_{p^n-1}$ , then there exists a finite subgroup  $F$  such that  $C_p \rtimes \mu_{n^2} = C_{n^2} \twoheadrightarrow \mu_n = C_n = C_{p-1} \cong \text{Aut}(C_p)$ .

If we want to calculate the Farrell-Tate cohomology of  $H^*(F, M)$  for  $p$ -complete module  $M$ , we look at the spectral sequence  $H^i(\mu_{n^2}, H^j(C_p, M))$ , then  $H^j(C_p, M)$  is  $p$ -complete as well, therefore  $H^k(F, M) = H^k(C_p, M)^{\mu_{n^2}}$ .

For example, if  $M = \mathbb{Z}_p$ ,  $\hat{H}^*(C_p, \mathbb{Z}_p) \cong \mathbb{Z}/p[\beta^{\pm 1}]$  where  $\beta \in \hat{H}^2(C_p, \mathbb{Z}_p)$ . As  $\beta^n$  is invariant,  $\hat{H}^*(F, \mathbb{Z}_p) \cong \mathbb{Z}/p[\beta^{\pm n}]$ .

35 Nov 13, 2023

Consider the split short exact sequence

$$1 \longrightarrow \mathbb{S}_n \longrightarrow \mathbb{G}_n \longrightarrow \text{Gal} \longrightarrow 1$$

then identifying  $\mathbb{S}_n = \mathcal{O}_n^\times$  gives a  $C_n = \text{Gal}(\mathbb{F}_{p^n}/\mathbb{F}_p)$ -action on it, where  $n = p-1$ . The maximal finite  $P$ -group in  $\mathbb{S}_n$  is isomorphic to  $C_p$ , and we identify  $\hat{H}^*(\mathbb{S}_n, -) \cong \hat{H}^*(N_{\mathbb{S}_n}(C_p), -)$  and similarly for  $\mathbb{G}_n$ . Recall we identify  $\mathcal{O}_n \left[ \frac{1}{p} \right]$  containing  $\mathbb{Q}_p(\xi_p)$  as the division algebra over  $\mathbb{Q}_p$ . Let  $C_{\mathbb{S}_n}$  be the centralizer of  $C_p$  in  $\mathbb{S}_n$ , then it is  $\mathbb{Q}_p(\xi_p) \cap \mathbb{S}_n = \mathbb{Z}_p[\xi_p]^\times$ , then this extends to a short exact sequence

$$\begin{array}{ccccccc} 1 & \longrightarrow & C_{\mathbb{S}_n} & \longrightarrow & N_{\mathbb{S}_n} & \longrightarrow & \text{Aut}(C_p) \cong C_n \longrightarrow 1 \\ & & \uparrow & & \uparrow & & \uparrow \\ & & \langle \tau^n \rangle & \longrightarrow & \langle \tau \rangle_{C_{n^2}} & \longrightarrow & C_n \end{array}$$

Let  $\omega$  be a primitive  $(p^n-1)$ -th root unity, which is contained in  $\mathbb{Z}_{p^n}^\times \hookrightarrow \mathbb{S}_n$ , then let  $\tau = \frac{p^n-1}{n^2}$ , then  $\tau$  is a primitive  $n^2$ -th root of unity. Therefore  $C_n$  is realized by conjugation of  $\tau$  modulo  $\tau^n$ . Correspondingly, we identify  $\mathbb{Z}_p[\xi_p]^\times$  to be  $C_n \times C_p \times \mathbb{Z}_p^n$  where  $C_n = \langle \tau^n \rangle$ . Now as an  $\text{Aut}(C_p)$ -module, we have  $\mathbb{Z}_p^n \cong \chi(0) \oplus \cdots \oplus \chi(n-1)$ , where  $\chi(k) \cong \mathbb{Z}_p$  but the generator acts on  $x$  by multiplication by  $\tau^{nk}$ .

Consider the spectral sequence

$$H^*(N/C_p, \hat{H}^*(C_p, M)) \Rightarrow \hat{H}^*(N_{\mathbb{S}_n}, M)$$

where  $M$  is  $p$ -complete. If  $H \triangleleft G$  and  $G/H$  has finite cohomological dimension, then there is a spectral sequence  $H^p(G/H, \hat{H}^q(H)) \Rightarrow \hat{H}^{p+q}(G)$  as well. With trivial coefficients, we have  $\hat{H}^*(C_p, \mathbb{Z}_p) \cong \mathbb{Z}/p\mathbb{Z}[\beta^{\pm 1}]$  for  $b \in H^2$ . Then  $C_{\mathbb{S}_n}$  acts trivially on  $b$  by  $\tau \cdot b = \tau^n b$ , where  $\tau^n$  is a  $(p-1)$ -th root of unity. Therefore  $H^*(C_{\mathbb{S}_n}/C_p, \hat{H}^*(C_p, \mathbb{Z}_p)) \cong$

$\mathbb{Z}/p\mathbb{Z}[b^{\pm 1}] \otimes \bigwedge(x_0, \dots, x_{n-1})$ , where  $C_{\mathbb{S}_n/C_p}$  as  $C_n \times \mathbb{Z}_p^n$  has trivial action on  $\hat{H}^*(C_p, \mathbb{Z}_p)$ . Similarly, we have  $\hat{H}^*(N_{\mathbb{S}_n/C_p}, \hat{H}^*(C_p, \mathbb{Z}_p)) \cong (\mathbb{Z}_p[b^{\pm 1}])^{\text{Aut}(C_p)}$ , where  $\tau \cdot x_i = (\tau^n)^{-i} x_i$ , so for  $|\beta| = 2n$  and  $y_i = 1 + 2i$ , then the cohomology is  $\mathbb{Z}/p\mathbb{Z}[\beta^{\pm 1}] \otimes \bigwedge(y_0, \dots, y_{n-1})$  where  $\beta = b^n$  and  $y_i = b^i x_i$ . One can calculate that both spectral sequences collapse (note  $H^*(C_{\mathbb{S}_n/C_p}, \hat{H}^*(C_p, \mathbb{Z}_p)) \Rightarrow H^*(C_{\mathbb{S}_n}, \mathbb{Z}_p)$  since  $C_{\mathbb{S}_n} = C_p \times C_{\mathbb{S}_n/C_p}$ ).

We now study over coefficients in  $(E_n)_* = (E_n)_0[u^{\pm 1}]$ , where  $(E_n)_0$  is the universal deformation ring for a formal group law where  $|u| = -2$ . In particular, there is an  $\mathbb{G}_n$ -action on this group. This makes  $(E_n)_0 \cong \mathbb{Z}_{p^n}[[u_1, \dots, u_{n-1}]] \supseteq \mathfrak{m} = (p, u_1, \dots, u_{n-1})$  modulo  $(p, \mathfrak{m}^2)$ . By Hopkins-Miller,  $\mathbb{Z}_{p^n}\{u, uu_1, \dots, uu_{n-1}\}$  is  $C_p$ -isomorphic to the reduced regulars  $\bar{P}_{C_p}$  modulo  $(p, \mathfrak{m}^2)$ .